American Heart Journal

The international publication for the study of the heart and circulation

Editor in Chief: **DEAN T. MASON, M.D.**

Associate Editor: **ANTHONY N. DeMARIA, M.D.**

Assistant Editors: **EZRA A. AMSTERDAM, M.D.**
NAJAM A. AWAN, M.D.
GARRETT LEE, M.D.
THOMAS A. RIEMENSCHNEIDER, M.D.
ZAKAUDDIN VERA, M.D.

Managing Editor: **LESLIE J. SILVERNAIL**

Editorial Board

W. Gerald Austen, M.D., *Boston*
Daniel S. Berman, M.D., *Los Angeles*
J. Thomas Bigger, M.D., *New York*
William J. Bommer, M.D., *Davis*
Lawrence I. Bonchek, M.D., *Milwaukee*
Lawrence S. Cohen, M.D., *New Haven*
Lawrence H. Cohn, M.D., *Boston*
John J. Collins, Jr., M.D., *Boston*
C. Richard Conti, M.D., *Gainesville*
Mary Allen Engle, M.D., *New York*
Harvey Feigenbaum, M.D., *Indianapolis*
James M. Foerster, M.D., *Davis*
Victor F. Froelicher, M.D., *San Diego*
Paul C. Gillette, M.D., *Houston*
John F. Goodwin, M.D., *London*
Antonio M. Gotto, Jr., M.D., *Houston*
Rolf M. Gunnar, M.D., *Maywood*
Fred J. Harris, M.D., *Davis*
W. Proctor Harvey, M.D., *Washington, D.C.*
Richard H. Helfant, M.D., *Philadelphia*
Walter L. Henry, M.D., *Irvine*
J. O'Neal Humphries, *Columbia*
J. Willis Hurst, M.D., *Atlanta*
Mark E. Josephson, M.D., *Philadelphia*
James A. Joye, M.D., *Davis*
William B. Kannel, M.D., *Framingham*
Norman M. Kaplan, M.D., *Dallas*
Samuel Kaplan, M.D., *Cincinnati*
Joseph A. Kisslo, M.D., *Durham*
Lawrence J. Laslett, M.D. *Davis*
Michael Lesch, M.D., *Chicago*
Floyd D. Loop, M.D., *Cleveland*
Reginald I. Low, M.D., *Davis*
Robert I. Levy, M.D., *Bethesda*
Dan G. McNamara, M.D., *Houston*
Richard R. Miller, M.D., *Houston*
Robert A. O'Rourke, M.D., *San Antonio*
Joseph K. Perloff, M.D., *Los Angeles*
Bertram Pitt, M.D., *Ann Arbor*
Charles E. Rackley, M.D., *Birmingham*
William C. Roberts, M.D., *Bethesda*
David J. Sahn, M.D., *Tucson*
Robert C. Schlant, M.D., *Atlanta*
Pravin M. Shah, M.D., *Los Angeles*
Walter Sommerville, M.D., *London*
James T. Willerson, M.D., *Dallas*
David O. Williams, M.D., *Providence*
Roger A. Winkle, M.D., *Stanford*
Robert F. Zelis, M.D., *Hershey*
Douglas P. Zipes, M.D., *Indianapolis*

VOLUME 104

JULY-DECEMBER 1982

VOLUME 104

COPYRIGHT© 1982 BY

THE C.V. MOSBY COMPANY

All rights reserved

Printed in the United States of America

Author index*

A

Abe H, 1382 (BC)
Abe I, 1351
Abe K, 1339
Abendroth R-R, 996
Aderka D, 320 (L)
Adomian GE, 1328
Aellig WH, 346
Ahmed SS, 869 (BC)
Aisner SC, 306 (BC)
Akyurekli Y, 1247
Aldrich RF, 816
Al-Khader A, 515
Allen HD, 156 (BC)
Alpert JS, 1119
Amery A, 174 (L), 1395
Amsterdam EA, 921, 1224
Anderson C, 1368
Anderson CM, 323 (L)
Andries F, 269
Aristimuno GG, 109
Arnett EA, 306 (BC)
Ashburn WL, 27, 238, 1027
Ashley WW, 185
Atkinson AJ, 203
Atwood E, 238
Aviram A, 884 (L)
Awan NA, 1211, 1125, 1127, 1224

B

BalaSubramanian V, 417
Baltch AL, 1335
Bannerman KS,1203
Bantea C, 1114 (L)
Barboza JS, 595
Barlow CH, 529
Barlow JB, 635 (BC)
Baron P, 1107 (BC)
Barrett MJ, 872 (BC)
Barron JV, 156 (BC)
Bateman T, 182
Battaglini JW, 816
Bauernfeind RA, 43, 185, 775
Beazell JW, 1328
Beder SD, 155 (BC)
Beeder C, 963
Beeson TP, 269
Belhassen B, 1117
Benchimol A, 974
Benito LP, 164 (BC)
Bennett CM, 421
Berger HJ, 1022
Berman DS, 182, 653, 709
Berman W Jr, 612
Bernheim J, 884 (L)
Bhakta RD, 1319
Bhargava V, 27, 100
Biegler TL, 1288
Bigger JT Jr, 746
Bircks W, 996
Bishop RL, 311 (BC)
Blackbourne BD, 312 (BC)
Blackman MS, 794
Blaser LE, 1114 (L)
Bloor CM, 158 (BC)
Blum J, 1004
Blum M, 884 (L)
Boelaert J, 174 (L)
Bolomo NJ, 1385 (BC)
Bonchek LI, 865
Borggrefe M, 996
Bornheimer JF, 1288
Bouchard RJ, 57
Bradof J, 1112 (BC)
Bragg-Remschel DA, 323 (L)
Braunwald E, 665
Bravo EL, 803
Breithardt G, 996
Brensike JF, 816
Browne KF, 163 (BC)
Bruce RA, 1032
Brundage BH, 879 (BC)
Bulkley BH, 561
Bulpitt C, 1395
Buonocore E, 628 (BC)
Bussey HI, 289

C

Cabin HS, 216, 306 (BC)
Caillé G, 1394 (L)
Cameron JS, 551
Camm AJ, 194
Campbell RW, 269
Campbell WB, 234
Canada WB, 874 (BC)
Cannon PJ, 92, 1203
Cannon RE, 438
Carabello BA, 136, 939
Caralis DG, 57
Carey PH, 254
Carliner NH, 456
Carr AA, 479
Carrasco G HA, 595
Carruth JE, 545
Carruthers SG, 451
Carter CH, 425
Cattaert A, 1395
Chacko KA, 885 (L)
Chang-Sing Yang D, 1377 (BC)
Charuzi Y, 963
Chatterjee K, 575, 1137
Chaudhuri TK, 145
Chee TP, 974
Chen T-S, 72
Cheng TO, 870 (BC)
Chilson DA, 163 (BC)
Chokshi AB, 1393 (L)
Chow MSS, 1016
Chun PKC, 160 (BC)
Ciro E, 643 (BC)
Clark BJ, 334
Cobb FR, 1232
Cody RJ, 1180, 1184, 1190
Coelho A, 166 (BC)
Cohen L, 881 (BC)
Cohn JN, 587, 725, 1159
Cohn K, 13
Colan SD, 175 (L)
Cole FE, 109
Coleman RE, 1232
Comerota A, 939
Conklin CM, 799
Conroy J, 1335
Conti CR, 173 (L), 226
Corday E, 740
Costello D, 100
Covit A, 1180
Cowley MJ, 946
Coxon R, 1164
Craig MWM, 417
Cramer M, 746
Crawford ES, 1271
Crawford MH, 145, 1054, 1061
Creager MA, 176 (L), 1147
Cutler BS, 66

D

Dahl C, 1244
Dalen JE, 1119
Daneels R, 174 (L)
Dansky R, 635 (BC)
Davies PTG, 173 (L)
Davies R, 1022
Davison R, 203
Dawson SL, 303 (BC)
Del Negro AA, 276
DeMaria AN, 874 (BC), 921
Denenberg BS, 939
Denes P, 169 (BC)
Dersham GH, 551
Désourdy F, 1394 (L)
Desser KB, 974
Detre KM, 816
DeWood MA, 521
Diamond GA, 732
DiBianco R, 77
DiCarlo LA Jr, 879 (BC)
Dicicco BS, 153 (BC), 303 (BC)
Dietrich P, 921
Dillon T, 612
Dittrich H, 100
Dório RJ, 1076
Dormandy J, 1364
Drayer JI, 660
Dreslinski GR, 109
Duncan WJ, 780
Dunlap TE, 1373 (BC)
Dymond DS, 977

E

Eddy H, 425
Egeblad H, 1105 (BC)
Eisemann G, 881 (BC)
Eldar M, 1357
Elkayam U, 20
Elliott AT, 977
Engel WK, 77
Epstein SE, 816, 1071
Ernst E, 1364
Escala EL, 1203
Ewy GA, 1244, 1303
Ezri MD, 169 (BC)

F

Fagard R, 174 (L), 1395
Famularo MA, 1086

*July, pp. 1-184; August, Part 1, pp. 185-332; August, Part 2, pp. 333-520; September, pp. 521-708; October, Part 1, pp. 709-890; October, Part 2, pp. 891-952; November, Part 1, pp. 953-1124; November, Part 2, pp. 1125-1228; December, pp. 1229-1430. *Key*: In this index, (BC) = brief communication, and (L) = letter to the Editor.

Fanchamps A, 388, 486
Farina M, 1045
Faxon DP, 176 (L), 1147
Feikes HL, 329
Feldman RL, 173 (L), 226
Fenster PE, 1244
Ferguson RK, 1117
Ferrufino O, 164 (BC)
Fievez M, 637 (BC)
Finerman WB Jr, 987
Fioretti P, 1109 (BC)
Firth BG, 234
Fisher JD, 177
Fisher ML, 456
Fisher MR, 816
Flessas AP, 311 (BC)
Fletcher RD, 77
Flute PT, 1364
Forrester JS, 521, 732
Fouad FM, 803, 827
Franciosa JA, 587
Francis GS, 725
Franken P, 637 (BC)
Freedom RM, 780
French WJ, 104
Freund GC, 912
Friday KJ, 323 (L)
Friedman HS, 72
Friedman MJ, 1303
Friedman PL, 769
Friesinger GC, 504
Frishman W, 496, 1154
Frithz G, 413
Fritz JK, 899
Froelicher V, 238
Frohlich ED, 109
Fuenmayor A, 595
Fuentes F, 912
Fukiyama K, 1351
Furmanski M, 1328

G

Gaeta JM, 912
Gaffney Nelson J, 1271
Gallant SL, 322 (L)
Gallastegui J, 603
Gang E, 746
Gangadharan V, 925
Garcia E, 653
Garner D, 104
Gault JH, 939
Gay JA, 77
Gelfand MJ, 785
Gentzler RD, 939
Georgiev M, 175 (L)
Gerber KH, 27
Gill CC, 628
Gillum RF, 839, 852
Ginsburg JH, 963
Glade LB, 109
Glassock RJ, 421
Glover MU, 1400
Gold HK, 946
Goldberg RJ, 57
Goldberg SJ, 156 (BC)
Goldberger AL, 617, 888 (L), 1381 (BC)
Goldhaber SZ, 305 (BC)
Goldsmith SR, 725
Goldstein DS, 1071
Goldstein JE, 834
Goldstein S, 649 (L)
Gomes GI, 1393 (L)
Gonasun LM, 374, 421, 438, 482, 496
Gopalaswamy C, 1104 (BC)
Gordon S, 925
Goto T, 1339
Gottdiener JS, 77, 312 (BC)
Gottschalk A, 1022
Gould KL, 912
Gould L, 1104 (BC), 1393 (L), 1377 (BC)
Grant CT, 852
Gratz E, 1071
Grauer K, 320 (L)
Gray RJ, 182, 799
Grenadier E, 156 (BC), 1384 (BC)
Griffin JC, 1004
Gross BW, 1032
Groves BM, 221
Gutgesell HP, 155 (BC), 1107 (BC)
Gutman J, 653

H

Haendchen RV, 740
Haffty BG, 311 (BC)
Haider B, 869 (BC)
Hall K, 646 (BC)
Halperin JL, 176 (L), 1147
Hamer A, 987
Hamilton BP, 421, 432, 456
Hamilton J, 432
Hamilton JH, 456
Hammer MC, 1335
Hammond JJ, 456
Han J, 551
Handler JB, 1400
Hanlon JT, 581
Hannah H III, 881 (BC)
Hansen JE, 442
Harken AH, 51, 529, 735, 1076
Harrigan PR, 875 (BC)
Harris L, 702
Harrison DC, 198, 209
Harston WE, 504
Haruyama T, 1339
Haselgrove J, 529
Hauser AM, 925
Hawkins E, 1107 (BC)
Hawley RJ, 77
Hecht HS, 567, 718
Heckman JL, 1309
Heger JJ, 163 (BC), 1095 (BC)
Hegesh J, 1376 (BC)
Heikkilä J, 464
Helfant RH, 697
Hellman DJ, 1303
Henis MMJ, 587
Hennekens CH, 305 (BC)
Henry WL, 20
Henuzet C, 637 (BC)
Henze E, 1, 953
Herfkens RL, 879 (BC)
Hermanovich J, 1211
Hernandez J, 515
Heupler FA, 690
Higgins CB, 27, 254
Hill JA, 226
Hillis LD, 234
Hiramatsu K, 1346
Hirsh PD, 234
Hirsh SA, 173 (L)
Hiszczynskyj R, 881 (BC)
Hiwatari M, 1339
Hockett D, 104
Holly AN, 575
Holman BL, 769
Honig HS, 115
Hooper WW, 1027
Hope RR, 537
Hopkins JM, 567, 718
Horowitz LN, 51, 753, 1076
Horwitz LD, 221
Hosler MH, 496
Hossack KF, 1032
Howard R, 641 (BC)
Howard T, 175 (L)
Hoy RH, 175 (L)
Huaghom F, 581
Huang MTC, 316 (BC)
Huang SK, 169 (BC)
Huber-Smith MJ, 1255
Huggenholtz PG, 1109 (BC)
Hunsaker JC, 303 (BC)

I

Ichimaru Y, 320 (L)
Imai Y, 1339
Inglessis G, 595
Inoue H, 1382 (BC)
Iseri LT, 20
Ishikawa H, 1262
Isner JM, 1278
Isomura S, 1262
Iverson N, 43
Iwasaki T, 1011
Izukawa T, 780

J

Jacks-Nagle VL, 456
Jackson GM, 158 (BC)
Janicki JS, 587
Janos GG, 785
Janoski AH, 456
Jarrott B, 407
Jee L, 606
Jensen D, 238
Joe P, 921
Johnson LL, 92
Johnson WD, 823
Johnson YE, 92
Johnston WD, 20
Jones RH, 85, 1232
Josephson MA, 567
Josephson ME, 51, 753, 1076, 1097 (BC)
Joswig BC, 1400

K

Kaba H, 649 (L)
Kabell G, 537
Kahl FR, 690
Kalischer A, 1203
Kanaide H, 1039
Kaplan S, 785
Karahalios SE, 718
Karbenn U, 996
Kasai Y, 1339
Kates RE, 198
Katz AM, 622
Katz NM, 312 (BC)
Kawano Y, 1351
Kayser KL, 649 (L), 823
Keefe DL, 198
Keidar S, 1384 (BC)
Keiser HR, 1071
Kelsey SE, 816
Kennedy HL, 57, 1086
Kennedy JW, 899
Kent KM, 276, 690
Keon WJ, 1247
Kerber RE, 263
Kerin NZ, 690
Kienzle MG, 753
Kim JS, 1288
Kim SG, 177
Kinasewitz GT, 587
Kindig JR, 269
Kinoshita A, 1262

Kirkendall WM, 333, 421, 432, 456
Kirsten E, 198
Kishel JC, 303 (BC)
Kitazume H, 648 (L)
Kocienski DE, 1027
Koilpillai C, 641 (BC)
Kondos GT, 603
Korecky B, 1247
Kostis JB, 36, 496
Kowey PR, 769
Krahn J, 646 (BC)
Kramer BL, 1172, 1197
Kramer JR, 648 (L)
Kramer W, 1294
Krikler DM, 702
Krohn BG, 1392 (L)
Krupp P, 486
Kuan P, 314 (BC)
Kubota T, 1346
Kugler J, 1154
Kuo PT, 36
Kuramoto K, 1011
Kusumi F, 1032
Kuwajima I, 1011

L

Lachman AS, 622
Laddu A, 874 (BC)
LaFoliette LE, 13
Lakin PC, 1112 (BC)
Laks MM, 104
Lam W, 603, 775
Langlois S, 473
Langrall H, 482
Laragh JH, 834, 1184
Latson TW, 221
Lazzara R, 537
Leach CN Jr, 622
Lee BY, 66
Lee G, 874 (BC), 921
Lee K, 921
Lee PK, 1016
Leijala MA, 885 (L)
Leinbach RC, 946
LeJemtel TH, 1154
Lerman BB, 1097 (BC)
Levine TB, 587, 1159
Levisman JA, 732
Levy PS, 43
Levy RI, 324, 816
Lewis KB, 306 (BC)
Liang C-s, 690
Lijnen P, 174 (L)
Likoff MJ, 877 (BC)
Lipton MJ, 879 (BC)
Litchfield R, 263
Livelli F, 746
Lloyd EA, 1095 (BC)
Loh IK, 816, 963
Loke J, 1022
Louis WJ, 407
Low RI, 874 (BC), 921
Lown B, 249, 491, 769
Lui H, 921, 1211, 1224
Luke JL, 303 (BC)
Lutgen J, 1319
Lynch PR, 1309

M

MacAlpin RN, 648 (L)
MacPhee AA, 109
Maddahi J, 182, 653
Malmud LS, 939
Mancini GBJ, 617
Mandel WJ, 799, 987
Mann RH, 939
Mann S, 417
Marcus FI, 1244
Maron BJ, 276, 643 (BC)
Marshall LA, 963
Marshall M, 92
Martin JL, 753
Martin RP, 136
Martinucci L, 1377 (BC)
Maskin CS, 1154
Mason DT, 874 (BC), 891, 921, 1127, 1211, 1224
Mason JW, 1004
Mason T, 603
Massie BM, 581, 1125, 1172, 1197
Mathew PK, 316 (BC)
Mathew TM, 316 (BC)
Matloff JM, 799
Matos JA, 177
Matrai A, 1364
Matsushita S, 1011
Matthay RA, 1022
Maurer AH, 939
Maurer G, 740
Maxwell M, 421
Mayers DL, 136
Mazer MS, 875 (BC)
McCabe CH, 1147
McCann DS, 1255
McCann WD, 939
McCarthy D, 877 (BC)
McGrath FJ, 276
McIntosh CL, 276, 306 (BC)
McKay CR, 575
McKirnan MD, 238
McManus BM, 153 (BC), 303 (BC), 312 (BC)
McNamara DG, 155 (BC), 1107 (BC)
McNeil JJ, 407
Meatherall B, 646 (BC)
Medina N, 1215
Meerbaum S, 740
Mehlman DG, 1378 (BC)
Meier J, 364
Meister SG, 872 (BC)
Melidossian CD, 1147
Meltzer RS, 1109 (BC)
Messer JV, 169 (BC)
Messerli FH, 109
Mickle D, 521
Mifune J, 1011
Milei J, 1385 (BC)
Miller RR, 1271
Milne JR, 194
Miyamoto A, 653
Molina C, 595
Moodie DS, 628
Moore RJ, 968
Moores WY, 1381 (BC)
Moreyra AE, 36
Mori KW, 1373 (BC)
Moroki N, 1011
Morrow AG, 276
Mortensen SA, 1105 (BC)
Moser KM, 1027
Motro M, 1376 (BC)
Muller JE, 672
Mulligan OF, 479
Mullins CE, 155 (BC)
Murakami M, 1011

N

Nacioglu O, 20
Nagaya T, 1262
Nakamura M, 1039
Nakano FS, 732
Nakashima Y, 827
Nanda NC, 308 (BC)
Needham KE, 1224
Nemerovski M, 709
Neufeld HM, 1357, 1376 (BC)
Niarchos AP, 834
Nicholls MG, 812
Nieminen MS, 464
Nikolic G, 311 (BC)
Noble RJ, 690
Nuñez L, 164 (BC)

O

Ockene IS, 1119
O'Connell JB, 1114 (L)
Ohkawa S, 1382 (BC)
Ohsugi J, 1262
Ohta T, 1262
Okada RD, 1303
Oliver JM, 164 (BC)
Olley PM, 780, 1045
Omae T, 1351
Opie LH, 606
O'Reilly DJ, 622
Oriel RJ, 762
Ormiston JA, 718
O'Rourke RA, 145, 1054, 1061
Orth R, 1381 (BC)
Ostermeyer J, 996
Otsuka K, 320 (L), 649 (L)

P

Packer M, 1215
Palant A, 1384 (BC)
Palileo EV, 43, 185
Palmeri ST, 1232
Pantano JA, 762
Parker M, 203
Parmley WW, 575, 1137
Passamani ER, 816
Patel S, 1393 (L)
Pavlicek W, 628 (BC)
Pedraza PM, 823
Peer G, 884 (L)
Penner B, 646 (BC)
Pepine CJ, 226
Pescarmona JE, 57
Peter CA, 85
Peter T, 521, 987
Peters FA, 209
Petropoulos AT, 185
Pfaff MJ, 732
Pichler M, 709
Pinkhas J, 320 (L)
Pitterman A, 1071
Plotnick GD, 456
Pocock WA, 635 (BC)
Podrid PJ, 491, 769
Polis O, 637 (BC)
Pollick C, 641 (BC)
Popat KD, 1373 (BC)
Porter JM, 66
Porta TA, 575
Povzhitkov M, 740
Powers ER, 1203
Prakash NS, 974
Prause JA, 521
Prichard BNC, 515
Proclemer A, 1097 (BC)
Proudfit WL, 648 (L)
Prystowsky EN, 163 (BC), 1095 (BC)

Q

Quinones MA, 1271

R

Rabinowitz B, 1357
Rabson JL, 646 (BC)
Raeder EA, 249
Raftery EB, 417
Rakowski H, 641 (BC)
Ramos RG, 925
Rand MJ, 407
Rangno RE, 473
Ratib O, 1, 953
Raymond M, 182
Reduto LA, 912
Reed KL, 1368
Rees GM, 977
Regan TJ, 869 (BC), 968
Reich T, 66
Reichle FA, 66
Reiffel JA, 746
Reison DS, 1203
Reynolds J, 238
Rich S, 603, 775
Richards KL, 145
Richardson JM, 816
Ritchie JL, 899
Ritterman JB, 1032
Roberts AJ, 329
Roberts CS, 638 (BC)
Roberts WC, 115, 153 (BC), 216, 276, 303 (BC), 306 (BC), 312 (BC), 638 (BC), 643 (BC), 1278
Robinson JD, 320 (L)
Robinson M, 456
Rodgers DM, 872 (BC)
Roeck W, 20
Rolnitzky L, 746
Rosen KM, 43, 166 (BC), 185, 775
Rosenberg KM, 939
Roskamm N, 512
Ross DL, 1004
Rotmensch HH, 1117
Rouleau J-L, 1137
Rowe RD, 780
Rowland E, 702
Rozanski A, 182, 653
Rubenfire M, 690
Rubin RN, 939
Ruffy R, 323 (L)
Rutledge J, 1224
Ryan TJ, 176 (L), 1147

S

Sahn DJ, 156 (BC), 1368
Saito H, 649 (L)
Sakai M, 1011
Sambasivan V, 1288
Samek L, 512
Sands MJ Jr, 622, 1112 (BC)
Santo M, 320 (L)
Sappington JB, 622
Sasse L, 651 (L)
Sato K, 1339
Sato M, 1339
Schaffer C, 1335
Schelbert HR, 1, 953
Scherlag BJ, 537
Schick EC Jr, 690, 1147
Schiller NB, 897 (BC)
Schlepper M, 1294
Schneeweiss A, 1376 (BC)
Schnugg SJ, 718
Schön H, 1, 953
Schuessler JS, 1271
Schurgers M, 174 (L)
Schuster AH, 308 (BC)
Schwartz DC, 785
Schwartz JB, 198
Schwarz HJ, 357
Sciacca RR, 92, 1203
Scogin JT, 66
Seibert A, 20
Seipel L, 996
Serruys PW, 1109 (BC)
Sethna DH, 799
Seto K, 320 (L), 649 (L)
Shaer G, 1180
Shah AA, 1271
Shah PK, 709
Shaher R, 1045
Shao-zhen Q, 1115 (L)
Shapiro W, 874 (BC)
Sharpe DN, 1164
Shawl FA, 160 (BC)
Shayegani M, 1335
Sheikh A, 603
Shell WE, 521
Shellock F, 709
Shem-Tov A, 1376 (BC)
Shenker L, 1368
Sheps DS, 1392 (L)
Sherrill LN, 479
Sherry S, 939
Sideris EB, 1045
Silverman ME, 545
Simson MB, 529
Singh BN, 567, 718
Singh JB, 311 (BC)
Slavin RG, 438
Slutsky RA, 27, 100, 254, 1027
Smalling RW, 912
Smith RP, 1335
Sobonya R, 1368
Sondheimer HM, 794
Sones FM Jr, 648 (L)
Sonnenblick EH, 1154
Sorensen SG, 145
Sorensen SM, 221
Sorkin RP, 1255, 1373 (BC)
Sotillo JF, 164 (BC)
Southworth W, 323 (L)
Spanarkel M, 1071
Spann JF, 939
Spear JF, 1076
Spodick DH, 311 (BC), 652 (L), 886 (L), 1392 (L)
Spooner E, 1045
Spragg RG, 1027
Spurrell RAJ, 194, 977
Staessen J, 1395
Stamm RB, 136
Starling MR, 1054
Steiger JF, 1255
Steingo L, 635 (BC)
Stephens JD, 977
Sterling R, 912
Stern R, 581
Stewart GH, 1309
Stone DL, 977
Stone J, 1203
Stone JM, 1319
Stone NJ, 816
Stone PH, 13, 672
Strandness DE, 66
Strasberg B, 43, 185, 166 (BC), 775
Stumpf DA, 887 (L)
Sturzenhofecker P, 512
Suarez DH, 109
Sue DY, 442
Sugiura M, 1382 (BC)
Sutphen NT, 1335
Swan HJC, 521, 709, 732
Swerdlow CD, 1004
Swiryn S, 43, 185, 775

T

Tabatznik B, 690
Taichman GC, 1247
Tajima J, 1339
Takatsu F, 1262
Takeda P, 921
Takeshita A, 1039
Takeya Y, 1351
Talano JV, 1378 (BC)
Tamari I, 1357
Tan KS, 1328
Tarazi RC, 803, 827, 886 (L)
Tavel ME, 269
Taylor H, 407
Terry RW, 681, 690
Thormann J, 1288
Thorsen NL, 496
Timmis GC, 925
Tobis J, 20
Tokarsky JM, 1255
Toltzis R, 1071
Topic N, 1172, 1197
Toyama J, 1262
Tse TF, 1016
Tubau J, 238
Tucker CR, 209
Turi ZG, 672

U

Ueda K, 1011, 1382 (BC)
Ueyama C, 1382 (BC)
Upton MT, 1232

V

Valdes-Cruz LM 156 (BC), 1368
van den Brand M, 1109 (BC)
Van Eeghem P, 174 (L)
Vanhees L, 1395
Vas R, 732
Ventura HO, 109
Verrier RL, 249
Verschueren LJ, 174 (L)
Vetrovec GW, 946
Vieweg WVR, 1400
Vilhelmsen R, 1105 (BC)
Virmani R, 1278
von Wichert P, 446

W

Waggoner AD, 1271
Walden RJ, 515
Walker W, 912
Wallace RB, 276, 312 (BC)
Waller BF, 153 (BC), 276, 303 (BC), 638 (BC)
Wallis DE, 1114 (L)
Walsh RA, 221, 1061
Wan-chun C, 1115 (L)
Wang RYC, 1016
Wanta-Matthews M, 912
Ward DE, 194
Warner HF, 939
Warren SE, 1400
Warren SG, 175 (L)
Wasserman K, 442
Waxman HL, 1097 (BC)
Webb Kavey R-E, 794
Weber KT, 587, 877 (BC)
Weber MA, 660
Wei JY, 561

Weinstein J, 496, 894
Weiss D, 987
Weiss MB, 1203
Weisse AB, 968
Weitzman RE, 421
Welch WJ, 166 (BC)
Welzel D, 512
West J, 587
Westveer DC, 925
Wetstein L, 529
White D, 606
Whiting JS, 732
Whitman HH, 834
Willerson JT, 234
Williams G, 1180
Winer N, 421, 425
Winkle RA, 209, 323 (L), 1004
Winsky M, 881 (BC)
Wisenberg G, 953
Witztum KR, 238, 1027
Wolf NM, 872 (BC)
Wolf PL, 158 (BC)
Wynne J, 769

X

Xin-xiang F, 1115 (L)

Y

Yabek SM, 612
Yamada K, 1262
Yamada T, 1346
Yamagishi F, 1346
Yanaga T, 320 (L), 649 (L)
Yang DC-S, 1104 (BC)
Yaroslavsky A, 884 (L)
Yeh HL, 996
Yiannikas J, 628
Yoshinaga K, 1339
Yu, DYC, 1016
Yu Y, 515
Yushak M, 1215

Z

Zaret BL, 1022
Zhen-jia P, 1115 (L)
Zielonka J, 769
Zipes DP, 163 (BC), 1095 (BC)
Ziskin MC, 1309
Zuckerman GL, 872 (BC)
Zweifler AJ, 812
Zweil P, 968

Subject index*

A

Ablation, cardioversion, and defibrillation: the sparkling joules of internal cardiac stimulation (Fisher, Matos, and Kim), 177

Acknowledgment to reviewers, 1229

Acoustical performance of the stethoscope: a comparative analysis (Kindig et al.), 269

Adenocarcinoma, rectal, cardiac metastasis of diagnosed by two-dimensional echocardiography (Henuzet et al.), 637 (Brief comm.)

Adrenergic-adrenoreceptor-blocking drugs, clinical pharmacology of (Louis et al.), 407

Adrenoreceptor-adrenergic-blocking drugs, clinical pharmacology of (Louis et al.), 407

Adverse reactions to pindolol administration (Gonasun and Langrall), 482

Aggregation, platelet, and plasma catecholamines, in vivo, in acute myocardial infarction (Sorkin et al.), 1255

Airway obstruction, reversibility of bronchospasm in (von Wichert), 446

reactivity and beta blockade in asthmatic patients, value of exercise in testing (Sue, Hansen, and Wasserman), 442

Alpha-receptor blockade, demonstration of additive benefits of: variable clinical response to long-term angiotensin inhibition in severe heart failure (Kugler et al.), 1154

Ambulatory ECG monitoring systems, ST segment performance of (Gallant), 322 (Letter)

Reply (Bragg-Remschel, Anderson, and Winkle), 323

electrocardiography in the assessment of pacemaker function (Famularo and Kennedy), 1086

American-born Caucasians versus Malayans, contrasting progression of mitral stenosis in (Joswig et al.), 1400

Amiodarone administration, sinoatrial arrest due to lidocaine injection in sick sinus syndrome during (Keidar, Grenadier, and Palant), 1384 (Brief comm.)

-aprindine interaction, possible (Southworth, Friday, and Ruffy), 323 (Lettter)

—benefits and risks in perspective (Rotmensch, Belhassen, and Ferguson), 1117

electrophysiologic observations of concealed ventricular depolarizations due to (Ezri et al.), 169 (Brief comm.)

therapy, chronic, electrophysiologic effects of in patients with ventricular arrhythmias (Finerman et al.), 987

Amplitude of atrial fibrillation waves in the geriatric population, clinicopathologic study on determinants of the (Inoue et al.), 1382 (Brief comm.)

Anemia, hemolytic, severe, after attempted repair of paraprosthetic mitral regurgitation (Goldberger, Orth, and Moores), 1381 (Brief comm.)

Aneurysm, left ventricular false (acquired cor triatriatum): complication of active infective endocarditis of the aortic valve with ring abscess treated by valve replacement (McManus et al.), 312 (Brief comm.)

septal, ventricular, paradoxically bulging into the left ventricular outflow tract, two-dimensional echocardiographic features of (Barron et al.), 156 (Brief comm.)

ventricular, noninvasive detection of by combined two-dimensional echocardiography and equilibrium radionuclide angiography (Sorensen et al.), 145

Aneurysmal formation or rupture, left ventricular incision midway between the mitral anulus and stumps of the papillary muscles during mitral valve excision with or without: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278

Aneurysmectomy, left ventricular, combined exercise radionuclide and hemodynamic evaluation of (Dymond et al.), 977

Angina pectoris and cold air exposure (Hirsh), 173 (Lettter)

Reply (Feldman and Conti), 173

chronic stable, advantages and disadvantages of combination therapy in: calcium-channel blockers and beta blockers (Krikler, Harris, and Rowland), 702

nifedipine therapy in: evaluation of safety and side effects (Terry), 681

refractory, efficacy of nifedipine therapy for (Stone, Turi, and Muller), 672

stable, randomized double-blind study of pindolol in patients with (Harston and Friesinger), 504

with pindolol, treatment of: the significance of intrinsic sympathomimetic activity of beta blockers (Kostis et al.), 496

Prinzmetal's, ischemic and postischemic ventricular wall motion abnormalities in provoked by hyperventilation (Egeblad, Vilhelmsen, and Mortensen), 1105 (Brief comm.)

recurrent, reinfarction, and reocclusion following intracoronary thrombolysis with streptokinase in acute myocardial infarction, importance of follow-up medical and surgical approaches to prevent (Lee et al.), 921

unstable, inpatient treatment of: clinical perspective and sequential management (Helfant), 697

sequential intracoronary streptokinase and transluminal angioplasty in with evolving myocardial infarction (Meltzer et al.), 1109 (Brief comm.)

variant, regional coronary artery dilation response in (Hill et al.), 226

Angiocardiography, radionuclide, resting and exercise, assessment of left ventricular function by, following acute myocardial infarction (Upton et al.), 1232

Angiographic and clinical results of intracoronary thrombolysis in syndromes of unstable ischemia (Vetrovec et al.), 946

and scintigraphic techniques, complementary roles of in assessment of the extent of coronary artery disease (Gutman et al.), 653

methods and dye dilution, ejection fraction derived using (French et al.), 104

Angiography, computer-enhanced digital: correlation of clinical assessment of left ventricular ejection fraction and regional wall motion (Vas et al.), 732

coronary, sequential, reliability of assessing change with (Detre et al.), 816

digital intravenous, analysis by: effects of acute incremental volume overload on cardiac chamber size, function, and the pulmonary circulation (Slutsky, Carey, and Higgins), 254

subtraction, in the evaluation of congenital abnormalities of the aorta and aortic arch (Moodie et al.), 628

subtraction, left ventricular imaging with using intravenous contrast injection and fluoroscopic exposure levels (Tobis et al.), 20

radionuclide, diagnosis of tricuspid regurgitation by (Gould, Yang, and Gopalaswamy), 1104 (Brief comm.)

postoperative evaluation of the Fontan procedure by (Janos et al.), 785

Angioplasty, transluminal, and sequential intracoronary streptokinase in unstable angina with evolving myocardial infarction (Meltzer et al.), 1109 (Brief comm.)

Angiotensin-converting enzyme inhibition with captopril, the special importance of: vasodilator therapy of severe congestive heart failure (Awan and Mason), 1127

inhibition in severe heart failure, long-term, variable clinical response to: demonstration of additive benefits of alpha-receptor blockade (Kugler et al.), 1154

Antianginal and antiarrhythmic activity, (pindolol's), evaluation of: pindolol postmyocardial infarction study (Welzel et al.), 512

*July, pp. 1-184; August, Part 1, pp. 185-332; August, Part 2, pp. 333-520; September, pp. 521-708; October, Part 1, pp. 709-890; October, Part 2, pp. 891-952; November, Part 1, 953-1124; November, Part 2, pp. 1125-1228; December, pp. 1229-1430.

Antiarrhythmic and antianginal activity, (pindolol's), evaluation of: pindolol postmyocardial infarction study (Welzel et al.), 512
drug(s), class I, prediction of response to during electrophysiologic study of ventricular tachycardia (Swiryn et al.), 43
efficacy, decreased incidence of at electrophysiologic study associated with the use of a third extrastimulus (Swerdlow et al.), 1004
surgery, effects of on late ventricular potentials recorded by precordial signal averaging in patients with ventricular tachycardia (Breithardt et al.), 996
Anticoagulation, embolization, prophylactic pacemaker, and long-term success, consideration of in cardioversion of atrial fibrillation (Mancini and Goldberger), 617
Antihypertensive agents which inhibit sympathetic activity: potentially adverse effects of combination treatment (Drayer and Weber), 660
effects of nifedipine combined with cardioselective beta-adrenergic receptor antagonism by atenolol (Opie, Jee, and White), 606
of pindolol (Gonasun), 374
Anulus, mitral, and the stumps of the papillary muscles, left ventricular incision midway between during mitral valve excision with or without rupture or aneurysmal formation: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
Aorta and aortic arch, intravenous digital subtraction angiography in the evaluation of congenital abnormalities of the (Moodie et al.), 628
Aortic arch and aorta, intravenous digital subtraction angiography in the evaluation of congenital abnormalities of the (Moodie et al.), 628
ostium, separate, of the left anterior descending and left circumflex coronary arteries from the left aortic sinus of Valsalva (absent left main coronary artery) (Dicicco et al.), 153 (Brief comm.).
regurgitation, chronic, cardiac response to exercise in patients with (Peter and Jones), 85
radionuclide angiographic evaluation of left ventricular function by resting ejection rate during the first third of systole in patients with (Johnson et al.), 92
stenosis, sudden death in monitored by ear densitographic pulse and ECG (Nikolic et al.), 311 (Brief comm.)
valves, normally functioning, previously confirmed, documented development of severe stenoses of (Roberts et al.), 306 (Brief comm.)
Aortitis, Takayasu's, etiologic aspects of coagulopathy in (Kanaide, Takeshita, and Nakamura), 1039
Apical hypertrophic nonobstructive cardiomyopathy (Steingo et al.), 635 (Brief comm.)
Apnea, sleep, influence of on nocturnal arrhythmias in ischemic heart disease (Otsuka et al.), 320 (Lettter)
Aprindine-amiodarone interaction, possible (Southworth, Friday, and Ruffy), 323 (Lettter)
AR-L 115, new cardiotonic agent, hemodynamic and myocardial energetic changes induced by in patients with coronary artery disease (Thormann, Kramer, and Schlepper), 1294
Arrhythmias, cardiac, in apparently normal well-trained runners, prevalence and nature of (Pantano and Oriel), 762
nocturnal, in ischemic heart disease, influence of sleep apnea on (Otsuka et al.), 320 (Lettter)
supraventricular, postoperative, and the role of prophylactic digoxin in cardiac surgery (Chee et al.), 974
ventricular, and regional myocardial blood flow following one-stage and two-stage coronary artery occlusion in anesthetized dogs (Kabell et al.), 537
comparison of two methods of analyzing frequency of (Shapiro et al.), 874 (Brief comm.)
electrophysiologic effects of chronic amiodarone therapy in patients with (Finerman et al.), 987
occurring late after surgery for congenital heart disease, phenytoin therapy for (Kavey, Blackman, and Sondheimer), 794
pindolol for (Podrid and Lown), 491
Arterial disease patients, occlusive, chronic, multicenter controlled double-blind trial with objective assessment of: pentoxifylline efficacy in the treatment of intermittent claudication (Porter et al.), 66
Arterial—*continued*
pressure, cardiac rhythm, and urinary catecholamines in borderline and established essential hypertension, diurnal variations of (Messerli et al.), 109
Arteriovenous fistula, coronary, two-dimensional echocardiographic features of (Rodgers et al.), 872 (Brief comm.)
Artery(ies) disease, coronary, and bovine milk-fat globule membrane (Davies), 173 (Letter)
complementary roles of scintigraphic and angiographic techniques in assessment of extent of (Gutman et al.), 653
dilation response, regional coronary, in variant angina (Hill et al.), 226
great, transposition of, corrected, two-dimensional echocardiographic identification of high anatomically inserted venous atrioventricular valve in the diagnosis of (Motro et al.), 1376 (Brief comm.)
ventricular diastolic pressure-volume relations after the intra-atrial baffle operation for (Sideris et al.), 1045
spasm, coronary, placebo-controlled study in patients with: randomized withdrawal from nifedipine (Schick et al.), 690
surgery, coronary, in diabetics: 261 consecutive patients followed four to seven years (Johnson, Pedraza, and Kayser), 823
Asthma, effect on of a new beta blocker, pindolol (Cannon, Slavin, and Gonasun), 438
Asthmatic patients, value of exercise in testing beta blockade and airway reactivity in (Sue, Hansen, and Wasserman), 442
Asymmetric and symmetric left ventricular hypertrophy, coexistence of in a family with hypertrophic cardiomyopathy (Ciro, Maron, and Roberts), 643 (Brief comm.)
septal hypertrophy on echocardiographic study, left ventricular thrombus demonstrating canalization and mimicking (Pollick et al.), 641 (Brief comm.)
Atenolol, cardioselective beta-adrenergic receptor antagonism by, combined with antihypertensive effects of nifedipine (Opie, Jee, and White), 606
Atherosclerotic narrowing, fatal, of the right main coronary artery: origin of the left anterior descending or left circumflex coronary artery from the right (the true "left-main equivalent") (Roberts, Waller, and Roberts), 638 (Brief comm.)
plaques, relation of healed transmural myocardial infarct size to length of survival after acute myocardial infarction, age at death, and amount and extent of coronary arterial narrowing by: analysis of 70 necropsy patients (Cabin and Roberts), 216
Atrial dissociation, transient, due to digitalis toxicity (Condorelli, Dato, and Sindoni), 1102 (Brief comm.)
fibrillation, cardioversion of: consideration of embolization, anticoagulation, prophylactic pacemaker, and long-term success (Mancini and Goldberger), 617
rapid, during hypercalcemia despite verapamil therapy (Santo, Aderka, and Pinkhas), 320 (Letter)
waves in the geriatric population, clinicopathologic study on determinants of the amplitude of (Inoue et al.), 1382 (Brief comm.)
inversion, isolated (Chacko), 885 (Letter)
Reply (Leijala), 885
myxoma, invasive: documentation of malignant potential of cardiac myxomas (Hannah et al.), 881 (Brief comm.)
tachycardia, automatic, surgical excision of: anatomic and electrophysiologic correlates (Josephson et al.), 1076
Atrioventricular bypass tracts, concealed, and nodoventricular bypass tracts, supraventricular tachycardia associated with (Lerman et al.), 1097 (Brief comm.)
disc valve malfunction, detection of by two-dimensional echocardiography (Mehlman and Talano), 1378 (Brief comm.)
valve, venous, high anatomically inserted, two-dimensional echocardiographic identification of in the diagnosis of corrected transposition of the great arteries (Motro et al.), 1376 (Brief comm.)
Automated blood pressure measurements during exercise testing, evaluation of (Hossack et al.), 1032
Automatic atrial tachycardia, surgical excision of: anatomic and electrophysiologic correlates (Josephson et al.), 1076

Autonomic nervous system on coronary blood flow during partial stenosis, influence of the (Raeder, Verrier, and Lown), 249
responses and postural hemodynamics in chronic heart failure, effect of captopril on (Cody), 1190

B

Bacteremia following dental cleaning in patients with and without penicillin prophylaxis (Baltch et al.), 1335
Baffle operation, intra-atrial, for transposition of the great arteries, ventricular diastolic pressure-volume relations after the (Sideris et al.), 1045
Benefits and risks of amiodarone, in perspective (Rotmensch, Belhassen, and Ferguson), 1117
Beta-adrenergic receptor antagonism by atenolol, cardioselective, antihypertensive effects of nifedipine combined with (Opie, Jee, and White), 606
-adrenoceptor blockers, secondary prevention with in postmyocardial infarction patients (Staessen et al.), 1395
-blocking agents: are pharmacologic differences relevant? (Clark), 334
other, pharmacokinetic comparison of pindolol with (Meier), 364
blockade, acute, cardiac safety of: intrinsic sympathomimetic activity is superior to beta-1 selectivity (Heikkilä and Nieminen), 464
and airway reactivity in asthmatic patients, value of exercise in testing (Sue, Hansen, and Wasserman), 442
low-dose, in the treatment of chronic cardiac failure (Weber, Likoff, and McCarthy), 877 (Brief comm.)
blocker(s), a new, pindolol, effect on asthma of (Cannon, Slavin, and Gonasun), 438
and calcium-channel blockers: advantages and disadvantages of combination therapy in chronic stable angina pectoris (Krikler, Harris, and Rowland), 702
significance of intrinsic sympathomimetic activity of: treatment of angina pectoris with pindolol (Kostis et al.), 496
-blocking drugs, withdrawal of (Walden et al.), 515
-1 selectivity, intrinsic sympathomimetic activity is superior to: cardiac safety of acute beta blockade (Heikkilä and Nieminen), 464
-stimulant, oral long-acting, salbutamol, an, in patients with congestive heart failure (Mifune et al.), 1011
Biplane and single-plane contrast analyses of right ventricular function and size, comparison of (Slutsky et al.), 100
Björk-Shiley prosthetic valve, mitral, malfunction of due to septal interference (Jackson, Wolf, and Bloor), 158
Black populations, coronary heart disease in. I. Mortality and morbidity (Gillum), 839
II. Risk factors (Gillum and Grant), 852
Block, bundle branch, after coronary bypass surgery (Bantea), 1114 (Lettter)
Reply (Wallis and O'Connell), 1114
heart, complete, experimental production of by electrocoagulation in the closed chest dog (Beazell et al.), 1328
left stellate ganglion, effects of drugs and on the long QT syndrome (Milne et al.), 194
Blockade, alpha-receptor, demonstration of additive benefits of: variable clinical response to long-term angiotensin inhibition in severe heart failure (Kugler et al.), 1154
beta, low-dose, in the treatment of chronic cardiac failure (Weber, Likoff, and McCarthy), 877 (Brief comm.)
Blockers, beta-adrenoceptor, secondary prevention with in postmyocardial infarction patients (Staessen et al.), 1395
and calcium-channel blockers: advantages and disadvantages of combination therapy in chronic stable angina pectoris (Krikler, Harris, and Rowland), 702
calcium-channel, and beta blockers: advantages and disadvantages of combination therapy in chronic stable angina pectoris (Krikler, Harris, and Rowland), 702
:pharmacologic considerations (Braunwald), 665
Blocking agents, beta-adrenoceptor: are pharmacologic differences relevant? (Clark), 334
Blood cardioplegia in coronary artery bypass graft surgery, harvesting and protection of the saphenous vein associated with early delivery of (Feikes and Roberts), 329
Blood—*continued*
flow, coronary, during partial stenosis, influence of the autonomic nervous system on (Raeder, Verrier, and Lown), 249
myocardial, regional, and ventricular arrhythmias following one-stage and two-stage coronary artery occlusion in anesthetized dogs (Kabell et al.), 537
pool studies, gated equilibrium, a new technique for determination of right ventricular stroke counts from: assessment of regurgitant fraction and right and left ventricular function at rest and during exercise (Henze et al.), 953
pressure and heart rate response to treadmill exercise, effect of quinidine on (Fenster et al.), 1244
and plasma norepinephrine, effects of pindolol and methyldopa on (Winer, Carter, and Eddy), 425
and plasma renin activity: pindolol effects on (Glassock et al.), 421
measurements, automated, evaluation of during exercise testing (Hossack et al.), 1032
pulse rate, and the renin-angiotensin-aldosterone system in patients with essential hypertension, acute effects of the calcium antagonist, nifedipine, on (Hiramatsu et al.), 1346
sampling site and technique, influence of on thromboxane concentrations in patients with ischemic heart disease (Hirsh et al.), 234
Body surface isopotential maps and left ventriculograms, correlation between in patients with old inferoposterior myocardial infarction (Ohta et al.), 1262
Bookshelf, 1404
Borderline hypertension, diminished finger volume pulse in: evidence for early structural vascular abnormality (Zweifler and Nicholls), 812
elevated plasma catecholamines without alteration in cardiovascular responsiveness in young men with (Kawano et al.), 1351
Bovine milk-fat globule membrane, coronary artery disease and (Davies), 173 (Lettter)
Bridge(s), myocardial, clinical significance of (MacAlpin), 648 (Lettter)
Reply (Kramer et al.), 648
or coronary tunnel, the case against clinical significance of: origin of the left main from the right coronary artery or from the right aortic sinus with intramyocardial tunneling to the left side of the heart via the ventricular septum (Roberts et al.), 303 (Brief comm.)
Bronchospasm in airway obstruction, reversibility of (von Wichert), 446
Bundle branch block after coronary bypass surgery (Bantea), 1114 (Letter)
Reply (Wallis and O'Connell), 1114
reentry, sustained ventricular tachycardia due to (Lloyd et al.), 1095 (Brief comm.)
Bypass, coronary artery, sternal dehiscence, pneumopericardial tamponade post- (Shawl and Chun), 160 (Brief comm.)
graft surgery, coronary artery, harvesting and protection of the saphenous vein associated with early delivery of blood cardioplegia in (Feikes and Roberts), 329
surgery, coronary, bundle branch block after (Bantea), 1114 (Letter)
Reply (Wallis and O'Connell), 1114
tracts, nodoventricular and concealed atrioventricular, supraventricular tachycardia associated with (Lerman et al.), 1097 (Brief comm.)

C

Calcium antagonist, nifedipine, acute effects of on blood pressure, pulse rate, and the renin-angiotensin-aldosterone system in patients with essential hypertension (Hiramatsu et al.), 1346
-channel blockers and beta blockers: advantages and disadvantages of combination therapy in chronic stable angina pectoris (Krikler, Harris, and Rowland), 702
:pharmacologic considerations (Braunwald), 665
Cameral fistula, silent left coronary artery: probable cause of myocardial ischemia (Ahmed, Haider, and Regan), 869 (Brief comm.)

Canalization, left ventricular thrombus demonstrating and mimicking asymmetric septal hypertrophy on echocardiographic study (Pollick et al.), 641 (Brief comm.)
Captopril, acute and long-term effects of on exercise cardiac performance and exercise capacity in congestive heart failure (Topic, Kramer, and Massie), 1172
on left and right ventricular volumes and function in chronic heart failure (Massie, Kramer, and Topic), 1197
and other vasodilator drugs, comparison of: contrasting hemodynamic responses in severe heart failure (Packer, Medina, and Yushak), 1215
and sodium nitroprusside, comparative analysis of the hemodynamic actions of in severe chronic congestive heart failure (Hermanovich et al.), 1211
and terbutaline therapy, complementary combined, in severe chronic congestive heart failure (Awan et al.), 1224
chronotropic property of in hypertensive patients, evaluation of the (Imai et al.), 1339
determinants of clinical response and survival in patients with congestive heart failure treated with (Creager et al.), 1147
effect of on renal, coronary, and systemic hemodynamics in patients with severe congestive heart failure (Powers et al.), 1203
hemodynamic effects of in pulmonary hypertension of collagen vascular disease (Niarchos et al.), 834
in chronic heart failure, hemodynamic effects of: efficacy of low-dose treatment and comparison with prazosin (Sharpe and Coxon), 1164
in congestive heart failure: improved left ventricular function with decreased metabolic cost (Chatterjee, Rouleau, and Parmley), 1137
inotropic properties of (Colan), 175 (Letter)
Reply (Faxon et al.), 176
pharmacokinetics and the acute hemodynamic and hormonal response in patients with severe chronic congestive heart failure (Cody et al.), 1180
special importance of angiotensin-converting enzyme inhibition with: vasodilator therapy of severe congestive heart failure (Awan and Mason), 1127
the effect of on postural hemodynamics and autonomic responses in chronic heart failure (Cody), 1190
therapy, long-term, in congestive heart failure: serial hemodynamics and echocardiographic changes (Nakashima, Fouad, and Tarazi), 827
use of to estimate renin-angiotensin-aldosterone activity in the pathophysiology of chronic heart failure (Cody and Laragh), 1184
Cardiac arrhythmias in apparently normal well-trained runners, prevalence and nature of (Pantano and Oriel), 762
chamber size, effects of acute incremental volume overload on and on function, and the pulmonary circulation: analysis by digital intravenous angiography (Slutsky, Carey, and Higgins), 254
complications of Wilson's disease, fatal (Kuan), 314 (Brief comm.)
failure, chronic, low-dose beta blockade in the treatment of (Weber, Likoff, and McCarthy), 877 (Brief comm.)
hydatid cyst diagnosed by two-dimensional echocardiography (Oliver et al.), 164 (Brief comm.)
metastasis of rectal adenocarcinoma diagnosed by two-dimensional echocardiography (Henuzet et al.), 637 (Brief comm.)
myxomas, documentation of malignant potential of: invasive atrial myxoma (Hannah et al.), 881 (Brief comm.)
performance evaluations, hemodynamic, and radionuclide ejection fraction: patterns of exercise response in patients with severe left ventricular dysfunction (Hecht et al.), 718
exercise, and exercise capacity in congestive heart failure, acute and long-term effects of captopril on (Topic, Kramer, and Massie), 1172
improvement in by oral long-acting theophylline in chronic obstructive pulmonary disease (Matthay et al.), 1022
response to exercise in patients with chronic aortic regurgitation (Peter and Jones), 85
rhythm, arterial pressure, and urinary catecholamines in borderline and established essential hypertension, diurnal variations of (Messerli et al.), 109
Cardiac—*continued*
safety of acute beta blockade: intrinsic sympathomimetic activity is superior to beta-1 selectivity (Heikkilä and Nieminen), 464
size and function, radionuclide angiographic assessment of: right and left ventricular response to subcutaneous terbutaline in patients with chronic obstructive pulmonary disease (Hooper et al.), 1027
stimulation, internal, sparkling joules of: cardioversion, defibrillation, and ablation (Fisher, Matos, and Kim), 177
surgery, effect of on ventricular septal motion: assessment by intraoperative echocardiography and cross-sectional two-dimensional echocardiography (Waggoner et al.), 1271
postoperative supraventricular arrhythmias and the role of prophylactic digoxin in (Chee et al.), 974
tumors in children, role of two-dimensional echocardiography in detection of: space-occupying lesions of the myocardium (Duncan et al.), 780
Cardiography, thoracic impedance (Krohn), 1392 (Letter)
(Spodick), 1392
Reply (Sheps), 1392
Cardiomyopathy, apical hypertrophic nonobstructive (Steingo et al.), 635 (Brief comm.)
congestive, progression from hypertrophic obstructive cardiomyopathy to in a child (Beder et al.), 155 (Brief comm.)
hypertrophic, coexistence of asymmetric and symmetric left ventricular hypertrophy in a family with (Ciro, Maron, and Roberts), 643 (Brief comm.)
metabolic, a: Friedreich's disease (Stumpf), 887
obstructive, hypertrophic, progression from to congestive cardiomyopathy in a child (Beder et al.), 155 (Brief comm.)
Cardioplegia, blood, in coronary artery bypass graft surgery, harvesting and protection of the saphenous vein associated with early delivery of (Feikes and Roberts), 329
Cardioselective beta-adrenergic receptor antagonism by atenolol, antihypertensive effects of nifedipine combined with (Opie, Jee, and White), 606
Cardiotonic agent, new, AR-L 115, hemodynamic and myocardial energetic changes induced by in patients with coronary artery disease (Thormann, Kramer, and Schlepper), 1294
Cardiotoxicity, digitalis, and the hypothalamus (Otsuka et al.), 649 (Letter)
Cardiovascular disease in the Marfan syndrome, spectrum of: a clinico-morphologic study of 18 necropsy patients and comparison to 151 previously reported necropsy patients (Roberts and Honig), 115
disorders during pregnancy, medical treatment of (Tamari et al.), 1357
effects of hemodialysis (Spodick), 886 (Letter)
Reply (Tarazi), 886
response, relation of to the hypotensive effect of metoprolol (Fouad, Tarazi, and Bravo), 803
responsiveness in young men with borderline hypertension, elevated plasma catecholamines without alteration in (Kawano et al.), 1351
Cardioversion, defibrillation, and ablation: the sparkling joules of internal cardiac stimulation (Fisher, Matos, and Kim), 177
of atrial fibrillation: consideration of embolization, anticoagulation, prophylactic pacemaker, and long-term success (Mancini and Goldberger), 617
transvenous, of recurrent ventricular tachycardia, additional observations on (Zipes et al.) 163 (Brief comm.)
Carotid sinus syncope as a manifestation of parotid tumors (Mathew, Huang, and Mathew), 316 (Brief comm.)
Catecholamines, plasma, and platelet aggregation, in vivo, in acute myocardial infarction (Sorkin et al.), 1255
elevated, without alteration in cardiovascular responsiveness in young men with borderline hypertension (Kawano et al.), 1351
Caucasians, American-born, versus Malayans, contrasting progression of mitral stenosis in (Joswig et al.), 1400
Chagas' disease, left ventricular cineangiography in: detection of early myocardial damage (Carrasco et al.), 595

Children, cardiac tumors in, role of two-dimensional echocardiography in detection of: space-occupying lesions of the myocardium (Duncan et al.), 780
electrophysiologic effects of propranolol on sinus node function in (Yabek, Berman, and Dillon), 612
Cholesterol and noncardiovascular mortality, consideration of (Levy), 324
Chronic heart failure, acute and long-term effects of captopril on left and right ventricular volumes and function in (Massie, Kramer, and Topic), 1197
effect of captopril on postural hemodynamics and autonomic responses in (Cody), 1190
use of captopril to estimate renin-angiotensin-aldosterone activity in the pathophysiology of (Cody and Laragh), 1184
obstructive pulmonary disease, improvement in cardiac performance by oral long-acting theophylline in (Matthay et al.), 1022
right and left ventricular response to subcutaneous terbutaline in patients with: radionuclide angiographic assessment of cardiac size and function (Hooper et al.), 1027
Chronotropic property of captopril in hypertensive patients, evaluation of the (Imai et al.), 1339
Cineangiography, left ventricular, in Chagas' disease: detection of early myocardial damage (Carrasco et al.), 595
Cinefluoroscopy, diagnostic value of in the evaluation of prosthetic heart valve dysfunction (Sands et al.), 622
Circulatory control mechanisms in vasodepressor syncope (Goldstein et al.), 1071
Claudication, intermittent, pentoxifylline efficacy in the treatment of: multicenter controlled double-blind trial with objective assessment of chronic occlusive arterial disease patients (Porter et al.), 66
Class I antiarrhythmic drugs, prediction of response to during electrophysiologic study of ventricular tachycardia (Swiryn et al.), 43
Clearance of quinidine during peritoneal dialysis (Hall et al.), 646 (Brief comm.)
Clinical pharmacology of adrenergic-adrenoreceptor-blocking drugs (Louis et al.), 407
of pindolol (Aellig), 346
practice, intracoronary streptokinase in (Timmis et al.), 925
response and survival in patients with congestive heart failure treated with captopril, determinants of (Creager et al.), 1147
significance of myocardial bridges (MacAlpin), 648 (Letter) Reply (Kramer et al.), 648
signs and hemodynamic state in the early hours of transmural myocardial infarction, comparison of (Shell et al.), 521
Clinicopathologic study on determinants of the amplitude of atrial fibrillation waves in the geriatric population (Inoue et al.), 1382 (Brief comm.)
Coagulopathy in Takayasu's aortitis, etiologic aspects of (Kanaide, Takeshita, and Nakamura), 1039
Cold air exposure and angina pectoris (Hirsh), 173 (Letter) Reply (Feldman and Conti), 173
Collagen vascular disease, hemodynamic effects of captopril in pulmonary hypertension of (Niarchos et al.), 834
Collateral(s), coronary, protection of myocardium by the compensatory mechanism of after total occlusion of major coronary arteries shown in patients with familial hypercholesterolemia (Kuo, Kostis, and Moreyra), 36
vessels, coronary, functional significance of in patients with coronary artery disease (Horwitz et al.), 221
Combination treatment, potentially adverse effects of: antihypertensive agents which inhibit sympathetic activity (Drayer and Weber), 660
Complete heart block, experimental production of by electrocoagulation in the closed chest dog (Beazell et al.), 1328
Computer-enhanced digital angiography: correlation of clinical assessment of left ventricular ejection fraction and regional wall motion (Vas et al.), 732
Computerized tomography and two-dimensional echocardiography, noninvasive detection of proximal pulmonary artery thrombosis by (DiCarlo et al.), 879 (Brief comm.)
Conduction, intracardiac, left ventricular relaxation, and mitral valve prolapse in myotonia atrophica: assessment by digitized echocardiography and noninvasive His bundle recording (Gottdiener et al.), 77
Congenital heart disease, phenytoin therapy for ventricular arrhythmias occurring late after surgery for (Kavey, Blackman, and Sondheimer), 794
Congestive cardiomyopathy, progression from hypertrophic obstructive cardiomyopathy to in a child (Beder et al.), 155 (Brief comm.)
heart failure, acute and long-term effects of captopril on exercise cardiac performance and exercise capacity in (Topic, Kramer, and Massie), 1172
captopril in: improved left ventricular function with decreased metabolic cost (Chatterjee, Rouleau, and Parmley), 1137
chronic, minoxidil therapy in: acute plus long-term hemodynamic and clinical study (McKay et al.), 575
severe, captopril pharmacokinetics and the acute hemodynamic and hormonal response in patients with (Cody et al.), 1180
comparative analysis of the hemodynamic actions of captopril and sodium nitroprusside in (Hermanovich et al.), 1211
complementary combined captopril and terbutaline therapy in (Awan et al.), 1224
therapy of (Awan and Massie), 1125
determinants of acute and long-term response to converting enzyme inhibitors in (Levine and Cohn), 1159
hemodynamic effects of salbutamol, an oral long-acting beta-stimulant, in patients with (Mifune et al.), 1011
long-term captopril therapy in: serial hemodynamics and echocardiographic changes (Nakashima, Fouad, and Tarazi), 827
prazosin therapy in (Gould et al.), 1393 (Letter)
refractory, beneficial hemodynamic effects of intravenous diazoxide in (Massie et al.), 581
relationship of exercise capacity to resting left ventricular performance and basal plasma norepinephrine levels in patients with (Francis, Goldsmith, and Cohn), 725
severe, effect of captopril on renal, coronary, and systemic hemodynamics in patients with (Powers et al.), 1203
vasodilator therapy of: the special importance of angiotensin-converting enzyme inhibition with captopril (Awan and Mason), 1127
treated with captopril, determinants of clinical response and survival in patients with (Creager et al.), 1147
Constrictive pericarditis, massive organized intrapericardial hematoma mimicking (Dunlap et al.), 1373 (Brief comm.)
Contrast analyses of right ventricular function and size, comparison of single-plane and biplane (Slutsky et al.), 100
Control mechanisms, circulatory, in vasodepressor syncope (Goldstein et al.), 1071
Converting enzyme inhibitors in congestive heart failure, determinants of acute and long-term response to (Levine and Cohn), 1159
Cor triatriatum, acquired (left ventricular false aneurysm): complication of active infective endocarditis of the aortic valve with ring abscess treated by valve replacement (McManus et al.), 312 (Brief comm.)
Coronary angiography, sequential, reliability of assessing change with (Detre et al.), 816
arterial narrowing by atherosclerotic plaques, relation of healed transmural myocardial infarct sizse to length of survival after acute myocardial infarction, age at death, and amount and extent of: analysis of 70 necropsy patients (Cabin and Roberts), 216
arteriovenous fistula, two-dimensional echocardiographic features of (Rodgers et al.), 872 (Brief comm.)
artery(ies) bypass graft surgery, harvesting and protection of the saphenous vein associated with early delivery of blood cardioplegia in (Feikes and Roberts), 329
cameral fistula, silent left: probable cause of myocardial ischemia (Ahmed, Haider, and Regan), 869 (Brief comm.)
dilation response, regional, in variant angina (Hill et al.), 226

Coronary artery—*continued*
disease and bovine milk-fat globule membrane (Davies), 173 (Letter)
complementary roles of scintigraphic and angiographic techniques in assessment of extent of (Gutman et al.), 653
functional significance of coronary collateral vessels in patients with (Horwitz et al.), 221
hemodynamic and myocardial energetic changes induced by the new cardiotonic agent, AR-L 115, in patients with (Thormann, Kramer, and Schlepper), 1294
left main, patterns of exercise treadmill test performance in patients with: detection dependent on left coronary dominance or coexistent dominant right coronary disease (Stone, LaFollette, and Cohn), 13
phase analysis of radionuclide ventriculograms for the detection of (Ratib et al.), 1
reproducibility of equilibrium radionuclide ventriculography in patients with: response of left ventricular ejection fraction and regional wall motion to supine bicycle exercise (Hecht et al), 567
fistula, left, to left ventricular fistula: demonstration of coronary steal phenomenon (Cheng), 870 (Brief comm.)
left anterior descending and left circumflex, separate aortic ostium of the from the left aortic sinus of Valsalva (absent left main coronary artery) (Dicicco et al.), 153 (Brief comm.)
ligation, experimental, effects of oxygen administration and alteration in arterial P_{CO_2} on ischemic myocardial changes following (Weisse et al.), 968
right main, fatal atherosclerotic narrowing of the: origin of the left anterior descending or left circumflex coronary artery from the right (the true "left-main equivalent") (Roberts, Waller, and Roberts), 638 (Brief comm.)
spasm, placebo-controlled study in patients with: randomized withdrawal from nifedipine (Schick et al.), 690
surgery in diabetics: 261 consecutive patients followed four to seven years (Johnson, Pedraza, and Kayser), 823
blood flow during partial stenosis, influence of the autonomic nervous system on (Raeder, Verrier, and Lown), 249
bypass surgery, bundle branch block after (Bantea), 1114 (Letter)
Reply (Wallis and O'Connell), 1114
collateral(s), protection of myocardium by the compensatory mechanism of after total occlusion of major coronary arteries shown in patients with familial hypercholesterolemia (Kuo, Kostis, and Moreyra), 36
vessels, functional significance of in patients with coronary artery disease (Horwitz et al.), 221
death, sudden, definition of (Kayser), 649 (Letter)
Reply (Goldstein), 649
disease patients, long-term results of endocardial resection for sustained ventricular tachycardia in (Josephson, Harken, and Horowitz), 51
transient appearance of Q waves in during exercise electrocardiography: consideration of mechanisms and clinical importance (Bateman et al.), 182
heart disease in black populations. I. Mortality and morbidity (Gillum), 839
II. Risk factors (Gillum and Grant), 852
renal, and systemic hemodynamics in patients with severe congestive heart failure, effect of captopril on (Powers et al.), 1203
spasm, coronary thrombosis, and myocardial infarction: a hypothesis concerning the pathophysiology of acute myocardial infarction (Dalen, Ockene, and Alpert), 1119
steal phenomenon, demonstration of: left coronary artery-to-left ventricular fistula (Cheng), 870 (Brief comm.)
Cross-sectional two-dimensional echocardiography and intraoperative echocardiography, assessment of effect of cardiac surgery on ventricular septal motion by (Waggoner et al.), 1271
Cycle length, paced, in normals and patients with sinoatrial dysfunction, sinus node recovery time related to (Reiffel et al.), 746
Cyst, cardiac hydatid, diagnosed by two-dimensional echocardiography (Oliver et al.), 164 (Brief comm.)

D

Death, sudden coronary, definition of (Kayser), 649 (Letter)
Reply (Goldstein), 649
in aortic stenosis monitored by ear densitographic pulse and ECG (Nikolic et al.), 311 (Brief comm.)
Defibrillation, cardioversion, and ablation: the sparkling joules of internal cardiac stimulation (Fisher, Matos, and Kim), 177
Deficiency, magnesium, ECG changes in early stage of (Chen et al.), 1115 (Letter)
Dehiscence, sternal, pneumopericardial tamponade postcoronary artery bypass (Shawl and Chun), 160 (Brief comm.)
Dental cleaning, bacteremia following in patients with and without penicillin prophylaxis (Baltch et al.), 1335
Depolarizations, ventricular, concealed, due to amiodarone, electrophysiologic observations of (Ezri et al.), 169 (Brief comm.)
Diabetics, coronary artery surgery in: 216 consecutive patients followed four to seven years (Johnson, Pedraza, and Kayser), 823
Dialysis, peritoneal, clearance of quinidine during (Hall et al.), 646 (Brief comm.)
Diastolic and systolic function, left ventricular, in hyperthyroidism (Friedman et al.), 1303
pressure, ventricular, and volume relations after the intra-atrial baffle operation for transposition of the great arteries (Sideris et al.), 1045
Diazoxide, intravenous, in refractory congestive heart failure, beneficial hemodynamic effects of (Massie et al.), 581
Digital intravenous angiography, analysis by: effects of acute incremental volume overload on cardiac chamber size, function, and the pulmonary circulation (Slutsky, Carey, and Higgins), 254
subtraction angiography, intravenous, in the evaluation of congenital abnormalities of the aorta and aortic arch (Moodie et al.), 628
left ventricular imaging with using intravenous contrast injection and fluoroscopic exposure levels (Tobis et al.), 20
Digitalis cardiotoxicity and the hypothalamus (Otsuka et al.), 649 (Letter)
glycosides, the influence of quinidine and other agents on (Bussey), 289
toxicity, transient atrial dissociation due to (Condorelli, Dato, and Sindoni), 1102 (Brief comm.)
Digoxin concentration, serum, after quinidine administration, use of control steady-state serum digoxin levels for predicting (Friedman and Chen), 72
levels, serum, control steady-state, use of for predicting serum digoxin concentration after quinidine administration (Friedman and Chen), 72
maintenance dosage in obesity (Grauer and Robinson), 320 (Letter)
prophylactic, the role of in cardiac surgery, and postoperative supraventricular arrhythmias (Chee et al.), 974
Dilation response, regional coronary artery, in variant angina (Hill et al.), 226
Disc valve malfunction, atrioventricular, detection of by two-dimensional echocardiography (Mehlman and Talano), 1378 (Brief comm.)
Disopyramide phosphate, use of radionuclide ventriculography for assessment of changes in myocardial performance induced by (Kowey et al.), 769
Dissociation, transient atrial, due to digitalis toxicity (Condorelli, Dato, and Sindoni), 1102 (Brief comm.)
Diurnal variations of cardiac rhythm, arterial pressure, and urinary catecholamines in borderline and established essential hypertension (Messerli et al.), 109
Dosage forms of pindolol, three, observations on (Carruthers), 451
Drug(s) efficacy, antiarrhythmic, decreased incidence of at electrophysiologic study associated with the use of a third extrastimulus (Swerdlow et al.), 1004
vasodilator, other, and captopril, comparison of: contrasting hemodynamic responses in severe heart failure (Packer, Medina, and Yushak), 1215
Dual chamber sequential pacing management of sinus node dysfunction: advantages over single-chamber pacing (Stone, Bhakta, and Lutgen), 1319

Dye dilution and angiographic methods, ejection fraction derived using (French et al.), 104
Dysfunction, occult myocardial, objective evidence of in patients with frequent ventricular ectopy without clinically apparent heart disease (Kennedy et al.), 57
regional ischemic left ventricular, detection and assessment of severity of by digital fluoroscopy (Gerber et al.), 27
renal, with sulphinpyrazone therapy (Boelaert et al.), 174 (Letter)
Reply (Warren et al.), 175
sinus node, dual chamber sequential pacing management of: advantages over single-chamber pacing (Stone, Bhakta, and Lutgen), 1319

E

Ear densitographic pulse and ECG, sudden death in aortic stenosis monitored by (Nikolic et al.), 311 (Brief comm.)
Echocardiographic and serial hemodynamic changes in long-term captopril therapy in congestive heart failure (Nakashima, Fouad, and Tarazi), 827
correlations, two-dimensional and M-mode: postoperative abnormalities of interventricular septal motion (Kerber and Litchfield), 263
features of coronary arteriovenous fistula, two-dimensional (Rodgers et al.), 872 (Brief comm.)
of left ventricular myxoma, M-mode and two-dimensional (Mazer and Harrigan), 875 (Brief comm.)
of ventricular septal aneurysm, two-dimensional, paradoxically bulging into the left ventricular outflow tract (Barron et al.), 156 (Brief comm.)
identification, two-dimensional, of high anatomically inserted venous atrioventricular valve in the diagnosis of corrected transposition of the great arteries (Motro et al.), 1376 (Brief comm.)
measurement, two-dimensional, of left ventricular ejection fraction: prospective analysis of what constitutes an adequate determination (Stamm et al.), 136
Echocardiography and systolic time intervals, relative sensitivity of for assessing acute positive inotropic interventions in normal human subjects (Walsh, Crawford, and O'Rourke), 1061
intraoperative, and cross-sectional two-dimensional echocardiography, assessment of effect of cardiac surgery on ventricular septal motion by (Waggoner et al.), 1271
M-mode, in left bundle branch block: significance of frontal plane QRS axis (Strasberg et al.), 775
two-dimensional, and equilibrium radionuclide angiography, noninvasive detection of ventricular aneurysm by combined (Sorensen et al.), 145
cardiac hydatid cyst diagnosed by (Oliver et al.), 164 (Brief comm.)
cross-sectional, and intraoperative echocardiography, assessment of effect of cardiac surgery on ventricular septal motion by (Waggoner et al.), 1271
detection by: pericardiocentesis induced intrapericardial thrombus (Schuster and Nanda), 308 (Brief comm.)
of atrioventricular disc valve malfunction by (Mehlman and Talano), 1378 (Brief comm.)
early diagnosis of nontransmural myocardial infarction by (Loh et al.), 963
left atrial, evaluation of changes in left ventricular end-diastolic pressure by (Haendchen et al.), 740
real-time, determination of left ventricular ejection fraction by visual estimation during (Rich et al.), 603
Ectopy, ventricular, frequent, without clinically apparent heart disease, objective evidence of occult myocardial dysfunction in patients with (Kennedy et al.), 57
Ejection fraction derived using dye dilution and angiographic methods (French et al.), 104
left ventricular, and regional wall motion, correlation of clinical assessment of: computer-enhanced digital angiography (Vas et al.), 732
response of to supine bicycle exercise: reproducibility of equilibrium radionuclide ventriculography in patients with coronary artery disease (Hecht et al.), 567
determination of by visual estimation during real-time two-dimensional echocardiography (Rich et al.), 603
two-dimensional echocardiographic measurement of: prospective analysis of what constitutes an adequate determination (Stamm et al.), 136
Ejection fraction—*continued*
radionuclide, and hemodynamic cardiac performance evaluations: patterns of exercise response in patients with severe left ventricular dysfunction (Hecht et al.), 718
rate, resting, radionuclide angiographic evaluation of left ventricular function by during the first third of systole in patients with chronic aortic regurgitation (Johnson et al.), 92
Electrocardiographic changes following endocardial resection for ventricular tachycardia (Kienzle et al.), 753
in early stage of magnesium deficiency (Chen et al.), 1115 (Letter)
monitoring systems, ambulatory, ST segment performance of (Gallant), 322 (Letter)
Reply (Bragg-Remschel, Anderson, and Winkle), 323
Electrocardiography, ambulatory, in the assessment of pacemaker function (Famularo and Kennedy), 1086
exercise, transient appearance of Q waves in coronary disease during: consideration of mechanisms and clinical importance (Bateman et al.), 182
Electrocoagulation, experimental production of complete heart block by in the closed chest dog (Beazell et al.), 1328
Electrophysiologic effects of chronic amiodarone therapy in patients with ventricular arrhythmias (Finerman et al.), 987
of propranolol on sinus node function in children (Yabek, Berman, and Dillon), 612
observations of concealed ventricular depolarizations due to amiodarone (Ezri et al.), 169 (Brief comm.)
study, decreased incidence of antiarrhythmic drug efficacy at associated with the use of a third extrastimulus (Swerdlow et al.), 1004
Elimination kinetics, verapamil, prolongation of during chronic oral administration (Schwartz et al.), 198
Embolism, pulmonary, time trends in hospital mortality and diagnosis of (Goldhaber and Hennekens), 305 (Brief comm.)
Embolization, anticoagulation, prophylactic pacemaker, and long-term success, consideration of in cardioversion of atrial fibrillation (Mancini and Goldberger), 617
Encainide, intravenous, acute hemodynamic effects of in patients with heart disease (Tucker et al.), 209
End-diastolic pressure, left ventricular, evaluation of changes in by left atrial two-dimensional echocardiography (Haendchen et al.), 740
Endocardial resection for sustained ventricular tachycardia in coronary disease patients, long-term results of (Josephson, Harken, and Horowitz), 51
for ventricular tachycardia, electrocardiographic changes following (Kienzle et al.), 753
Endocarditis, active infective, of the aortic valve with ring abscess treated by valve replacement, complication of: acquired cor triatriatum (left ventricular false aneurysm) (McManus et al.), 312 (Brief comm.)
Enzyme inhibition, angiotensin-converting, with captopril, the special importance of: vasodilator therapy of severe congestive heart failure (Awan and Mason), 1127
inhibitors, converting, in congestive heart failure, determinants of acute and long-term response to (Levine and Cohn), 1159
Epinephrine, effects of on the electrophysiologic properties of Purkinje fibers surviving myocardial infarction (Cameron, Dersham, and Han), 551
Equilibrium radionuclide ventriculography, reproducibility of in patients with coronary artery disease: response of left ventricular ejection fraction and regional wall motion to supine bicycle exercise (Hecht et al.), 567
Essential hypertension, acute effects of the calcium antagonist, nifedipine, on blood pressure, pulse rate, and the renin-angiotensin-aldosterone system in patients with (Hiramatsu et al.), 1346
Etiologic aspects of coagulopathy in Takayasu's aortitis (Kanaide, Takeshita, and Nakamura), 1039
Exercise capacity and exercise cardiac performance in congestive heart failure, acute and long-term effects of captopril on (Topic, Kramer, and Massie), 1172
relationship of to resting left ventricular performance and basal plasma norepinephrine levels in patients with congestive heart failure (Francis, Goldsmith, and Cohn), 725

Exercise—*continued*
cardiac performance and exercise capacity in congestive heart failure, acute and long-term effects of captopril on (Topic, Kramer, and Massie), 1172
response to in patients with chronic aortic regurgitation (Peter and Jones), 85
electrocardiography, transient appearance of Q waves in coronary disease during: consideration of mechanisms and clinical importance (Bateman et al.), 182
in testing beta blockade and airway reactivity in asthmatic patients, value of (Sue, Hansen, and Wasserman), 442
protocols predischarge, selected treadmill, and six weeks post-infarction, superiority of for detecting ischemic abnormalities (Starling, Crawford, and O'Rourke), 1054
provocable right ventricular outflow tract tachycardia (Palileo et al.), 185
radionuclide and hemodynamic evaluation, combined, of left ventricular aneurysmectomy (Dymond et al.), 977
response in patients with severe left ventricular dysfunction, patterns of: radionuclide ejection fraction and hemodynamic cardiac performance evaluations (Hecht et al.), 718
supine and upright, in mitral stenosis, effects of nitroglycerin on (Bornheimer et al.), 1288
testing, evaluation of automated blood pressure measurements during (Hossack et al.), 1032
training, noninvasive assessment of changes in myocardial perfusion and ventricular performance following (Tubau et al.), 238
treadmill, effect of quinidine on the heart rate and blood pressure response to (Fenster et al.), 1244
test performance, patterns of in patients with left main coronary artery disease: detection dependent on left coronary dominance or coexistent dominant right coronary disease (Stone, LaFollette, and Cohn), 13
Exposure, cold air, and angina pectoris (Hirsh), 173 (Letter)
Reply (Feldman and Conti), 173
Extrastimulus, third, decreased incidence of antiarrhythmic drug efficacy at electrophysiologic study associated with the use of a (Swerdlow et al.), 1004

F

Failure, cardiac, chronic, low-dose beta blockade in the treatment of (Weber, Likoff, and McCarthy), 877 (Brief comm.)
Fatal atherosclerotic narrowing of the right main coronary artery: origin of the left anterior descending or left circumflex coronary artery from the right (the true "left-main equivalent") (Roberts, Waller, and Roberts), 638 (Brief comm.)
cardiac complications of Wilson's disease (Kuan), 314 (Brief comm.)
Fevers, hemorrhagic, viral, myocardial damage in (Milei and Bolomo), 1385 (Brief comm.)
Fibrillation, atrial, cardioversion of: consideration of embolization, anticoagulation, prophylactic pacemaker, and long-term success (Mancini and Goldberger), 617
rapid, during hypercalcemia despite verapamil therapy (Santo, Aderka, and Pinkhas), 320 (Letter)
ventricular, complicating acute myocardial infarction: reasons against the routine use of lidocaine (Carruth and Silverman), 545
waves, atrial, in the geriatric population, clinicopathologic study on determinants of the amplitude of (Inoue et al.), 1382 (Brief comm.)
Finger volume pulse, diminished, in borderline hypertension: evidence for early structural vascular abnormality (Zweifler and Nicholls), 812
First and second heart sounds, frequency analysis approach to the origin of the (Heckman et al.), 1309
Fistula, cameral, silent left coronary artery: probable cause of myocardial ischemia (Ahmed, Haider, and Regan), 869 (Brief comm.)
coronary arteriovenous, two-dimensional echocardiographic features of (Rodgers et al.), 872 (Brief comm.)
left coronary artery, to left ventricular fistula: demonstration of coronary steal phenomenon (Cheng), 870 (Brief comm.)
Fluoroscopy, digital, detection and assessment of severity of regional ischemic left ventricular dysfunction by (Gerber et al.), 27
Follow-up medical and surgical approaches, importance of to prevent reinfarction, reocclusion, and recurrent angina following intracoronary thrombolysis with streptokinase in acute myocardial infarction (Lee et al.), 921
Fontan procedure, postoperative evaluation of the by radionuclide angiography (Janos et al.), 785
Frequency analysis approach to the origin of the first and second heart sounds (Heckman et al.), 1309
of ventricular arrhythmias, comparison of two methods of analyzing (Shaipro et al.), 874 (Brief comm.)
Friedreich's disease: a metabolic cardiomyopathy (Stumpf), 887
Frontal plane QRS axis, significance of: M-mode echocardiography in left bundle branch block (Strasberg et al.), 775

G

Gated equilibrium blood pool studies, a new technique for determination of right ventricular stroke counts from: assessment of regurgitant fraction and right and left ventricular function at rest and during exercise (Henze et al.), 953
Geriatric population, clincopathologic study on determinants of the amplitude of atrial fibrillation waves in the (Inoue et al.), 1382 (Brief comm.)
Global ischemia and reperfusion, acute, myocardial responses to (Taichman et al.), 1247
Glycosides, digitalis, the influence of quinidine and other agents on (Bussey), 289
Great arteries, transposition of, corrected, two-dimensional echocardiographic identification of high anatomically inserted venous atrioventricular valve in the diagnosis of (Motro et al.), 1376 (Brief comm.)

H

Harvesting and protection of the saphenous vein associated with early delivery of blood cardioplegia in coronary artery bypass graft surgery (Feikes and Roberts), 329
Heart block, complete, experimental production of by electrocoagulation in the closed chest dog (Beazell et al.), 1328
disease, acute hemodynamic effects of intravenous encainide in patients with (Tucker et al.), 209
congenital, phenytoin therapy for ventricular arrhythmias occurring late after surgery for (Kavey, Blackman, and Sondheimer), 794
coronary, in black populations. I. Mortality and morbidity (Gillum), 839
II. Risk factors (Gillum and Grant), 852
ischemic, influence of blood sampling site and technique on thromboxane concentrations in patients with (Hirsh et al.), 234
failure, chronic, acute and long-term effects of captopril on left and right ventricular volumes and function in (Massie, Kramer, and Topic), 1197
effect of captopril on postural hemodynamics and autonomic responses in (Cody), 1190
hemodynamic effects of captopril in: efficacy of low-dose treatment and comparison with prazosin (Sharpe and Coxon), 1164
hydralazine in the long-term treatment of: lack of difference from placebo (Franciosa et al.), 587
use of captopril to estimate renin-angiotensin-aldosterone activity in the pathophysiology of (Cody and Laragh), 1184
congestive, acute and long-term effects of captopril on exercise cardiac performance and exercise capacity in (Topic, Kramer, and Massie), 1172
captopril in: improved left ventricular function with decreased metabolic cost (Chatterjee, Rouleau, and Parmley), 1137
chronic, minoxidil therapy in: acute plus long-term hemodynamic and clinical study (McKay et al.), 575
severe, captopril pharmacokinetics and the acute hemodynamic and hormonal response in patients with (Cody et al.), 1180

Heart failure, congestive, chronic, severe—*continued*
comparative analysis of the hemodynamic actions of captopril and sodium nitroprusside in (Hermanovich et al.), 1211
complementary combined captopril and terbutaline therapy in (Awan et al.), 1224
therapy of (Awan and Massie), 1125
determinants of acute and long-term response to converting enzyme inhibitors in (Levine and Cohn), 1159
hemodynamic effects of salbutamol, an oral long-acting beta-stimulant, in patients with (Mifune et al.), 1011
long-term captopril therapy in: serial hemodynamics and echocardiographic changes (Nakashima, Fouad, and Tarazi), 827
prazosin therapy in (Gould et al.), 1393 (Letter)
refractory, beneficial hemodynamic effects of intravenous diazoxide in (Massie et al.), 581
severe, effect of captopril on renal, coronary, and systemic hemodynamics in patients with (Powers et al.), 1203
vasodilator therapy of: the special importance of angiotensin-converting enzyme inhibition with captopril (Awan and Mason), 1127
treated with captopril, determinants of clinical response and survival in patients with (Creager et al.), 1147
severe, beneficial hemodynamic effects of intravenous terbutaline in patients with (Wang et al.), 1016
contrasting hemodynamic responses in: comparison of captopril and other vasodilator drugs (Packer, Medina, and Yushak), 1215
long-term angiotensin inhibition in, variable clinical response to: demonstration of additive benefits of alpha-receptor blockade (Kugler et al.), 1154
rate and blood pressure response to treadmill exercise, effect of quinidine on (Fenster et al.), 1244
sounds, first and second, frequency analysis approach to the origin of the (Heckman et al.), 1309
valve dysfunction, prosthetic, diagnostic value of cinefluoroscopy in the evaluation of (Sands et al.), 622
Hematoma, intrapericardial, massive organized, mimicking constrictive pericarditis (Dunlap et al.), 1373 (Brief comm.)
Hemodialysis, cardiovascular effects of (Spodick), 886 (Letter)
Reply (Tarazi), 886
Hemodynamic(s) actions of captopril and sodium nitroprusside in severe chronic congestive heart failure, comparative analysis of the (Hermanovich et al.), 1211
and exercise radionuclide evaluation, combined, of left ventricular aneurysmectomy (Dymond et al.), 977
and hormonal response, acute, in patients with severe chronic congestive heart failure, captopril pharmacokinetics and the (Cody et al.), 1180
and myocardial energetic changes induced by the new cardiotonic agent, AR-L 115, in patients with coronary artery disease (Thormann, Kramer, and Schlepper), 1294
cardiac performance evaluations and radionuclide ejection fraction: patterns of exercise response in patients with severe left ventricular dysfunction (Hecht et al.), 718
effects, beneficial, of intravenous diazoxide in refractory congestive heart failure (Massie et al.), 581
of captopril in chronic heart failure: efficacy of low-dose treatment and comparison with prazosin (Sharpe and Coxon), 1164
in pulmonary hypertension of collagen vascular disease (Niarchos et al.), 834
of intravenous encainide in patients with heart disease, acute (Tucker et al.), 209
terbutaline, beneficial, in patients with severe heart failure (Wang et al.), 1016
of salbutamol, an oral long-acting beta-stimulant, in patients with congestive heart failure (Mifune et al.), 1011
postural, and autonomic responses in chronic heart failure, effect of captopril on (Cody), 1190
renal, coronary, and systemic, effect of captopril on in patients with severe congestive heart failure (Powers et al.), 1203
responses in severe heart failure, contrasting: comparison of captopril and other vasodilator drugs (Packer, Medina, and Yushak), 1215
Hemodynamic—*continued*
serial, and echocardiographic changes in long-term captopril therapy in congestive heart failure (Nakashima, Fouad, and Tarazi), 827
state and clinical signs in the early hours of transmural myocardial infarction, comparison of (Shell et al.), 521
Hemolytic anemia, severe, after attempted repair of paraprosthetic mitral regurgitation (Goldberger, Orth, and Moores), 1381 (Brief comm.)
Hemorrhagic fevers, viral, myocardial damage in (Milei and Bolomo), 1385 (Brief comm.)
Hemorrheologic changes following acute myocardial infarction (Dormandy et al.), 1364
High-dose, brief intravenous streptokinase early in acute myocardial infarction (Spann et al.), 939
Hoechst-Roussel registry, efficacy and safety data from 209 United States cases in the: treatment of myocardial infarction with intracoronary streptokinase (Weinstein), 894
Hormonal and hemodynamic response, acute, in patients with severe chronic congestive heart failure, captopril pharmacokinetics and the (Cody et al.), 1180
Hyaluronidase, mechanism of action of in decreasing myocardial ischemia post coronary occlusion in the isolated perfused rabbit heart (Wetstein et al.), 529
Hydatid cyst, cardiac, diagnosed by two-dimensional echocardiography (Oliver et al.), 164 (Brief comm.)
Hydralazine in the long-term treatment of chronic heart failure: lack of difference from placebo (Franciosa et al.), 587
Hydrops fetalis, prenatal ultrasound diagnosis of hypoplastic left heart syndrome in utero associated with (Sahn et al.), 1368
Hyperacute T waves revisited (Goldberger), 888
Hypercalcemia, rapid atrial fibrillation during despite verapamil therapy (Santo, Aderka, and Pinkhas), 320 (Letter)
Hypercholesterolemia, familial, patients with, protection of myocardium by the compensatory mechanism of coronary collaterals after total occlusion of major coronary arteries shown in (Kuo, Kostis, and Moreyra), 36
Hypertension, borderline, diminished finger volume pulse in: evidence for early structural vascular abnormality (Zweifler and Nicholls), 812
elevated plasma catecholamines without alteration in cardiovascular responsiveness in young men with (Kawano et al.), 1351
essential, acute effects of the calcium antagonist, nifedipine, on blood pressure, pulse rate, and the renin-angiotensin-aldosterone system in patients with (Hiramatsu et al.), 1346
borderline and established, diurnal variations of cardiac rhythm, arterial pressure, and urinary catecholamines in (Messerli et al.), 109
once-daily pindolol in: an ambulatory assessment (Raftery et al.), 417
pindolol and systolic time intervals in patients with (Hammond et al.), 456
versus methyldopa for: comparison of adverse reactions (Carr, Mulligan, and Sherrill), 479
pulmonary, of collagen vascular disease, hemodynamic effects of captopril in (Niarchos et al.), 834
registry, new primary pulmonary (Blaser), 1114 (Letter)
therapeutic trials of pindolol in: comparison and combination with other drugs (Fanchamps), 388
with pindolol, once-a-day treatment of (Frithz), 413
Hypertensive patients, evaluation of the chronotropic property of captopril in (Imai et al.), 1339
treated with pindolol, pulmonary function in: a report of two studies (Hamilton, Hamilton, and Kirkendall), 432
Hyperthyroidism, left ventricular systolic and diastolic function in (Friedman et al.), 1303
Hypertrophic cardiomyopathy, coexistence of asymmetric and symmetric left ventricular hypertrophy in a family with (Ciro, Maron, and Roberts), 643 (Brief comm.)
nonobstructive cardiomyopathy, apical (Steingo et al.), 635 (Brief comm.)
obstructive cardiomyopathy, progression from to congestive cardiomyopathy in a child (Beder et al.), 155 (Brief comm.)

Hypertrophy, asymmetric and symmetric left ventricular, coexistence of in a family with hypertrophic cardiomyopathy (Ciro, Maron, and Roberts), 643 (Brief comm.)
septal, on echocardiographic study, left ventricular thrombus demonstrating canalization and mimicking (Pollick et al.), 641 (Brief comm.)
Hyperventilation, ischemic and postischemic ventricular wall motion abnormalities in Prinzmetal's angina provoked by (Egeblad, Vilhelmsen, and Mortensen), 1105 (Brief comm.)
Hypoplastic left heart syndrome in utero associated with hydrops fetalis, prenatal ultrasound diagnosis of (Sahn et al.), 1368
Hypotensive effect of metoprolol, relation of cardiovascular response to the (Fouad, Tarazi, and Bravo), 803
Hypothalamus and digitalis cardiotoxicity, the (Otsuka et al.), 649 (Letter)

I

Imaging, left ventricular, with digital subtraction angiography using intravenous contrast injection and fluoroscopic exposure levels (Tobis et al.), 20
Impedance cardiography, thoracic (Krohn), 1392 (Letter)
(Spodick), 1392
Reply (Sheps), 1392
In vivo platelet aggregation and plasma catecholamines in acute myocardial infarction (Sorkin et al.), 1255
Incremental volume overload, acute, effects of on cardiac chamber size, function, and the pulmonary circulation: analysis by digital intravenous angiography (Slutsky, Carey, and Higgins), 254
Infarction, myocardial, acute, hemorrheologic changes following (Dormandy et al.), 1364
serum phosphate shift in (Yaroslavsky et al.), 884 (Letter)
ventricular fibrillation complicating: reasons against the routine use of lidocaine (Carruth and Silverman), 545
before age 36 years in women: predominance of apparent nonatherosclerotic events (Wei and Bulkley), 561
effects of epinephrine on the electrophysiologic properties of Purkinje fibers surviving (Cameron, Dersham, and Han), 551
transmural, comparison of clinical signs and hemodynamic state in the early hours of (Shell et al.), 521
patients, post-myocardial, secondary prevention with beta-adrenoceptor blockers in (Staessen et al.), 1395
study, pindolol postmyocardial: evaluation of the drug's antiarrhythmic and antianginal activity (Welzel et al.), 512
Infective endocarditis, active, of the aortic valve with ring abscess treated by valve replacement, complication of: acquired cor triatriatum (left ventricular false aneurysm) (McManus et al.), 312 (Brief comm.)
Inferoposterior myocardial infarction, old, correlation between body surface isopotential maps and left ventriculograms in patients with (Ohta et al.), 1262
Infradiaphragmatic total anomalous pulmonary venous connection in siblings (Baron et al.), 1107 (Brief comm.)
Inotropic interventions, acute positive, in normal human subjects, relative sensitivity of echocardiography and systolic time intervals for assessing (Walsh, Crawford, and O'Rourke), 1061
properties of captopril (Colan), 175 (Letter)
Reply (Faxon et al.), 176
Inpatient treatment of unstable angina: clinical perspective and sequential management (Helfant), 697
Interventricular septal motion, postoperative abnormalities of: two-dimensional and M-mode echocardiographic correlations (Kerber and Litchfield), 263
Intra-atrial baffle operation for transposition of the great arteries, ventricular diastolic pressure-volume relations after the (Sideris et al.), 1045
Intracoronary streptokinase in clinical practice (Timmis et al.), 925
Intracoronary streptokinase—*continued*
sequential, and transluminal angioplasty in unstable angina with evolving myocardial infarction (Meltzer et al.), 1109 (Brief comm.)
thrombolysis in syndromes of unstable ischemia: angiographic and clinical results (Vetrovec et al.), 946
up to 18 hours after onset of pain in evolving myocardial infarction (Smalling et al.), 912
with streptokinase in acute myocardial infarction, importance of follow-up medical and surgical approaches to prevent reinfarction, reocclusion, and recurrent angina following (Lee et al.), 921
Intraoperative echocardiography and cross-sectional two-dimensional echocardiography, assessment of effect of cardiac surgery on ventricular septal motion by (Waggoner et al.), 1271
Intrapericardial hematoma, massive organized, mimicking constrictive pericarditis (Dunlap et al.), 1373 (Brief comm.)
thrombus, pericardiocentesis induced: detection by two-dimensional echocardiography (Schuster and Nanda), 308 (Brief comm.)
Intravenous digital subtraction angiography in the evaluation of congenital abnormalities of the aorta and aortic arch (Moodie et al.), 628
encainide, acute hemodynamic effects of in patients with heart disease (Tucker et al.), 209
streptokinase, high-dose, brief, early in acute myocardial infarction (Spann et al.), 939
terbutaline, beneficial hemodynamic effects of in patients with severe heart failure (Wang et al.), 1016
verapamil, role of in supraventricular tachyarrhythmias after open-heart surgery (Gray et al.), 799
Introduction to International Symposium on pindolol (Kirkendall), 333
Invasive atrial myxoma: documentation of malignant potential of cardiac myxomas (Hannah et al.), 881 (Brief comm.)
Inversion, isolated atrial (Chacko), 885 (Letter)
Reply (Leijala), 885
Ischemia, global, acute, and reperfusion, myocardial responses to (Taichman et al.), 1247
myocardial, mechanism of action of hyaluronidase in decreasing, post coronary occlusion in the isolated perfused rabbit heart (Wetstein et al.), 529
unstable, intracoronary thrombolysis in syndromes of: angiographic and clinical results (Vetrovec et al.), 946
Ischemic abnormalities, superiority of selected treadmill exercise protocols predischarge and six weeks postinfarction for detecting (Starling, Crawford, and O'Rourke), 1054
and postischemic ventricular wall motion abnormalities in Prinzmetal's angina provoked by hyperventilation (Egeblad, Vilhelmsen, and Mortensen), 1105 (Brief comm.)
heart disease, influence of blood sampling site and technique on thromboxane concentrations in patients with (Hirsh et al.), 234
of sleep apnea on nocturnal arrhythmias in (Otsuka et al.), 320 (Letter)
left ventricular dysfunction, regional, by digital fluoroscopy, detection and assessment of severity of (Gerber et al.), 27
myocardial changes following experimental coronary artery ligation, effects of oxygen administration and alteration in arterial P_{CO_2} on (Weisse et al.), 968
Isolated atrial inversion (Chacko), 885 (Letter)
Reply (Leijala), 885
Isopotential maps, body surface, and left ventriculograms, correlation between in patients with old inferoposterior myocardial infarction (Ohta et al.), 1262

L

Left and right ventricular function following first acute transmural myocardial infarction, radionuclide assessment of sequential changes in (Nemerovski et al.), 709
volumes and function in chronic heart failure, acute and long-term effects of captopril on (Massie, Kramer, and Topic), 1197

Left—*continued*
anterior descending coronary artery or left circumflex coronary artery from the right (the true "left-main equivalent"), origin of the: fatal atherosclerotic narrowing of the right main coronary artery (Roberts, Waller, and Roberts), 638 (Brief comm.)
atrial two-dimensional echocardiography, evaluation of changes in left ventricular end-diastolic pressure by (Haendchen et al.), 740
bundle branch block, M-mode echocardiography in: significance of frontal plane QRS axis (Strasberg et al.), 775
circumflex coronary artery or left anterior descending coronary artery from the right (the true "left-main equivalent"), origin of the: fatal atherosclerotic narrowing of the right main coronary artery (Roberts, Waller, and Roberts), 638 (Brief comm.)
coronary artery cameral fistula, silent: probable cause of myocardial ischemia (Ahmed, Haider, and Regan), 869 (Brief comm.)
-to-left ventricular fistula: demonstration of coronary steal phenomenon (Cheng), 870 (Brief comm.)
heart syndrome, hypoplastic, in utero associated with hydrops fetalis, prenatal ultrasound diagnosis of (Sahn et al.), 1368
main coronary artery, absent: separate aortic ostium of the left anterior descending and left circumflex coronary arteries from the left aortic sinus of Valsalva (Dicicco et al.), 153 (Brief comm.)
disease, patterns of exercise treadmill test performance in patients with: detection dependent on left coronary dominance or coexistent dominant right coronary disease (Stone, LaFollette, and Cohn), 13
origin of the from the right coronary artery or from the right aortic sinus with intramyocardial tunneling to the left side of the heart via the ventricular septum: the case against clinical significance of myocardial bridge or coronary tunnel (Roberts et al.), 303 (Brief comm.)
stellate ganglion block, effects of drugs and on the long QT syndrome (Milne et al.), 194
ventricular aneurysmectomy, combined exercise radionuclide and hemodynamic evaluation of (Dymond et al.), 977
cineangiography in Chagas' disease: detection of early myocardial damage (Carrasco et al.), 595
dysfunction, regional ischemic, detection and assessment of severity of by digital fluoroscopy (Gerber et al.), 27
severe, patterns of exercise response in patients with: radionuclide ejection fraction and hemodynamic cardiac performance evaluations (Hecht et al.), 718
ejection fraction, and regional wall motion, correlation of clinical assessment of: computer-enhanced digital angiography (Vas et al.), 732
determination of by visual estimation during real-time two-dimensional echocardiography (Rich et al.), 603
two-dimensional echocardiographic measurement of: prospective analysis of what constitutes an adequate determination (Stamm et al.), 136
end-diastolic pressure, evaluation of changes in by left atrial two-dimensional echocardiography (Haendchen et al.), 740
function, assessment of by resting and exercise radionuclide angiocardiography following acute myocardial infarction (Upton et al.), 1232
improved, with decreased metabolic cost: captopril in congestive heart failure (Chatterjee, Rouleau, and Parmley), 1137
radionuclide angiographic evaluation of by resting ejection rate during the first third of systole in patients with chronic aortic regurgitation (Johnson et al.), 92
imaging with digital subtraction angiography using intravenous contrast injection and fluoroscopic exposure levels (Tobis et al.), 20
incision midway between the mitral anulus and the stumps of the papillary muscles during mitral valve excision with or without rupture or aneurysmal formation: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
myxoma: M-mode and two-dimensional echocardiographic features (Mazer and Harrigan), 875 (Brief comm.)
Left ventricular—*continued*
performance, resting, and basal plasma norepinephrine levels in patients with congestive heart failure, relationship of exercise capacity to (Francis, Goldsmith, and Cohn), 725
potentials recorded by precordial signal averaging in patients with ventricular tachycardia, effects of antiarrhythmic surgery on (Briethardt et al.), 996
pseudoaneurysm, radionuclide diagnosis of (Gould, Yang, and Martinucci), 1377 (Brief comm.)
relaxation, mitral valve prolapse, and intracardiac conduction in myotonia atrophica: assessment by digitized echocardiography and noninvasive His bundle recording (Gottdiener et al.), 77
systolic and diastolic function in hyperthyroidism (Friedman et al.), 1303
thrombus demonstrating canalization and mimicking asymmetric septal hypertrophy on echocardiographic study (Pollick et al.), 641 (Brief comm.)
-to-left coronary artery fistula: demonstration of coronary steal phenomenon (Cheng), 870 (Brief comm.)
ventriculograms and body surface isopotential maps, correlation between in patients with old inferoposterior myocardial infarction (Ohta et al.), 1262
Lesions of the myocardium, space-occupying: role of two-dimensional echocardiography in detection of cardiac tumors in children (Duncan et al.), 780
Lidocaine injection, sinoatrial arrest due to in sick sinus syndrome during amiodarone administration (Keidar, Grenadier, and Palant), 1384 (Brief comm.)
levels, excessive serum, during maintenance infusions: mechanisms and prevention (Davison, Parker, and Atkinson), 203
reasons against the routine use of: ventricular fibrillation complicating acute myocardial infarction (Carruth and Silverman), 545
Ligation, coronary artery, experimental, effects of oxygen administration and alteration in arterial Pco_2 on ischemic myocardial changes following (Weisse et al.), 968
Long QT syndrome, the: effects of drugs and left stellate ganglion block (Milne et al.), 194
Low-dose beta blockade in the treatment of chronic cardiac failure (Weber, Likoff, and McCarthy) 877 (Brief comm.)

M

Macroreentrant ventricular tachycardia, sustained (Welch et al.), 166 (Brief comm.)
Magnesium deficiency, ECG changes in early stage of (Chen et al.), 1115 (Letter)
Malayans versus American-born Caucasians, contrasting progression of mitral stenosis in (Joswig et al.), 1400
Malignant potential of cardiac myxomas, documentation of: invasive atrial myxoma (Hannah et al.), 881 (Brief comm.)
Marfan syndrome, cardiovascular disease in the, spectrum of: a clinico-morphologic study of 18 necropsy patients and comparison to 151 previously reported necropsy patients (Roberts and Honig), 115
Massive organized intrapericardial hematoma mimicking constrictive pericarditis (Dunlap et al.), 1373 (Brief comm.)
Medical treatment of cardiovascular disorders during pregnancy (Tamari et al.), 1357
Metabolic cardiomyopathy, a: Friedreich's disease (Stumpf), 887
Metastasis, cardiac, of rectal adenocarcinoma diagnosed by two-dimensional echocardiography (Henuzet et al.), 637 (Brief comm.)
Methyldopa and pindolol on blood pressure and plasma norepinephrine, effects of (Winer, Carter, and Eddy), 425
versus pindolol for hypertension: comparison of adverse reactions (Carr, Mulligan, and Sherrill), 479
Metoprolol, hypotensive effect of, relation of cardiovascular response to the (Fouad, Tarazi, and Bravo), 803
propranolol, and pindolol, comparison of withdrawal phenomena after (Rangno and Langlois), 473

Milk-fat globule membrane, bovine, coronary artery disease and (Davies), 173 (Letter)
Minoxidil therapy in chronic congestive heart failure: acute plus long-term hemodynamic and clinical study (McKay et al.), 575
Mitral anulus and the stumps of the papillary muscles, left ventricular incision midway between during mitral valve excision with or without rupture or aneurysmal formation: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
regurgitation, clinically isolated, severe, chronic, pure, etiology of: analysis of 97 patients over 30 years of age having mitral valve replacement (Waller et al.), 276
paraprosthetic, severe hemolytic anemia after attempted repair of (Goldberger, Orth, and Moores), 1381 (Brief comm.)
stenosis, effects of nitroglycerin on supine and upright exercise in (Bornheimer et al.), 1288
in Malayans versus American-born Caucasians, contrasting progression of (Joswig et al.), 1400
valve disease, correction of without valve replacement (Bonchek), 865
excision with or without rupture or aneurysmal formation, left ventricular incision midway between the mitral anulus and stumps of the papillary muscles during: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
replacement, analysis of 97 patients over 30 years of age having: etiology of clinically isolated, severe, chronic, pure mitral regurgitation (Waller et al.), 276
M-mode and two-dimensional echocardiographic correlations: postoperative abnormalities of interventricular septal motion (Kerber and Litchfield), 263
features of left ventricular myxoma (Mazer and Harrigan), 875 (Brief comm.)
echocardiography in left bundle branch block: significance of frontal plane QRS axis (Strasberg et al.), 775
Monitoring systems, ambulatory ECG, ST segment performance of (Gallant), 322 (Letter)
Reply (Bragg-Remschel, Anderson, and Winkle), 323
Mortality and morbidity of coronary heart disease in black populations. Part I of study (Gillum), 839
hospital, time trends in and diagnosis of pulmonary embolism (Goldhaber and Hennekens), 305 (Brief comm.)
noncardiovascular, and cholesterol, consideration of (Levy), 324
Myocardial blood flow, regional, and ventricular arrhythmias following one-stage and two-stage coronary artery occlusion in anesthetized dogs (Kabell et al.), 537
bridges, clinical significance of (MacAlpin), 648 (Letter)
Reply (Kramer et al.), 648
damage, early, detection of with left ventricular cineangiography in Chagas' disease (Carrasco et al.) 595
in viral hemorrhagic fevers (Milei and Bolomo), 1385 (Brief comm.)
energetic and hemodynamic changes induced by the new cardiotonic agent, AR-L 115, in patients with coronary artery disease (Thormann, Kramer, and Schlepper), 1294
infarct size, transmural, healed, relation of to length of survival after acute myocardial infarction, age at death, and amount and extent of coronary arterial narrowing by atherosclerotic plaques: analysis of 70 necropsy patients (Cabin and Roberts), 216
infarction, acute, a hypothesis concerning the pathophysiology of: coronary spasm, coronary thrombosis, and myocardial infarction (Dalen, Ockene, and Alpert), 1119
assessment of left ventricular function by resting and exercise radionuclide angiocardiography following (Upton et al.), 1232
hemorrheologic changes following (Dormandy et al.), 1364
high-dose, brief intravenous streptokinase early in (Spann et al.), 939
importance of follow-up medical and surgical approaches to prevent reinfarction, reocclusion, and recurrent angina following intracoronary thrombolysis with streptokinase in (Lee et al.), 921
in vivo platelet aggregation and plasma catecholamines in (Sorkin et al.), 1255
Myocardial infarction, acute—*continued*
relation of healed transmural myocardial infarct size to length of survival after, age at death, and amount and extent of coronary arterial narrowing by atherosclerotic plaques: analysis of 70 necropsy patients (Cabin and Roberts), 216
serum phosphate shift in (Yaroslavsky et al.), 884 (Letter)
streptokinase in: Western Washington randomized trial-protocol and progress report (Kennedy, Fritz, Ritchie, et al.), 899
ventricular fibrillation complicating: reasons against the routine use of lidocaine (Carruth and Silverman), 545
before age 36 years in women: predominance of apparent nonatherosclerotic events (Wei and Bulkley), 561
coronary spasm, and coronary thrombosis: a hypothesis concerning the pathophysiology of acute myocardial infarction (Dalen, Ockene, and Alpert), 1119
effects of epinephrine on the electrophysiologic properties of Purkinje fibers surviving (Cameron, Dersham, and Han), 551
evolving, beneficial effects of intracoronary thrombolysis up to 18 hours after onset of pain in (Smalling et al.), 912
sequential intracoronary streptokinase and transluminal angioplasty in unstable angina with (Meltzer et al.), 1109 (Brief comm.)
inferoposterior, old, correlation between body surface isopotential maps and left ventriculograms in patients with (Ohta et al.), 1262
nontransmural, early diagnosis of by two-dimensional echocardiography (Loh et al.), 963
transmural, comparison of clinical signs and hemodynamic state in the early hours of (Shell et al.), 521
first acute, radionuclide assessment of sequential changes in left and right ventricular function following (Nemerovski et al.), 709
with intracoronary streptokinase, treatment of: efficacy and safety data from 209 United States cases in the Hoechst-Roussel registry (Weinstein), 894
ischemia, mechanism of action of hyaluronidase in decreasing, post coronary occlusion in the isolated perfused rabbit heart (Wetstein et al.), 529
silent left coronary artery cameral fistula, probable cause of (Ahmed, Haider, and Regan), 869 (Brief comm.)
performance induced by disopyramide phosphate, use of radionuclide ventriculography for assessment of changes in (Kowey et al.), 769
perfusion and ventricular performance following exercise training, noninvasive assessment of changes in (Tubau et al.), 238
responses to acute global ischemia and reperfusion (Taichman et al.), 1247
Myocardium, protection of by the compensatory mechanism of coronary collaterals after total occlusion of major coronary arteries shown in patients with familial hypercholesterolemia (Kuo, Kostis, and Moreyra), 36
space-occupying lesions of the: role of two-dimensional echocardiography in detection of cardiac tumors in children (Duncan et al.), 780
Myotonia atrophica, left ventricular relaxation, mitral valve prolapse, and intracardiac conduction in: assessment by digitized echocardiography and noninvasive His bundle recording (Gottdiener et al.), 77
Myxoma(s), cardiac, documentation of malignant potential of: invasive atrial myxoma (Hannah et al.), 881 (Brief comm.)
invasive atrial: documentation of malignant potential of cardiac myxomas (Hannah et al.), 881 (Brief comm.)
left ventricular: M-mode and two-dimensional echocardiographic features (Mazer and Harrigan), 875 (Brief comm.)

N

Nadolol dosage after propranolol (Tremblay et al.), 1394 (Letter)

Nervous system, autonomic, on coronary blood flow during partial stenosis, influence of the (Raeder, Verrier, and Lown), 249
Nifedipine, calcium antagonist, acute effects of on blood pressure, pulse rate, and the renin-angiotensin-aldosterone system in patients with essential hypertension (Hiramatsu et al.), 1346
combined with cardioselective beta-adrenergic receptor antagonism by atenolol, antihypertensive effects of (Opie, Jee, and White), 606
therapy for refractory angina pectoris, efficacy of (Stone, Turi, and Muller), 672
in angina pectoris: evaluation of safety and side effects (Terry), 681
withdrawal from, randomized: placebo-controlled study in patients with coronary artery spasm (Schick et al.), 690
Nitroglycerin, effects of on supine and upright exercise in mitral stenosis (Bornheimer et al.), 1288
Nocturnal arrhythmias in ischemic heart disease, influence of sleep apnea on (Otsuka et al.), 320 (Letter)
Nodoventricular and concealed atrioventricular bypass tracts, supraventricular tachycardia associated with (Lerman et al.), 1097 (Brief comm.)
Nonatherosclerotic events, predominance of apparent: myocardial infarction before age 36 years in women (Wei and Bulkley), 561
Noncardiovascular mortality and cholesterol, consideration of (Levy), 324
Noninvasive assessment of changes in myocardial perfusion and ventricular performance following exercise training (Tubau et al.), 238
detection of proximal pulmonary artery thrombosis by two-dimensional echocardiography and computerized tomography (DiCarlo et al.), 879 (Brief comm.)
Nontransmural myocardial infarction by two-dimensional echocardiography, early diagnosis of (Loh et al.), 963
Norepinephrine levels, basal plasma, and resting left ventricular performance in patients with congestive heart failure, relationship of exercise capacity to (Francis, Goldsmith, and Cohn), 725
plasma, and blood pressure, effects of pindolol and methyldopa on (Winer, Carter, and Eddy), 425

O

Obesity, digoxin maintenance dosage in (Grauer and Robinson), 320 (Letter)
Obstructive pulmonary disease, chronic, improvement in cardiac performance by oral long-acting theophylline in (Matthay et al.), 1022
Occlusive arterial disease patients, chronic, multicenter controlled double-blind trial with objective assessment of: pentoxifylline efficacy in the treatment of intermittent claudication (Porter et al.), 66
Occult myocardial dysfunction, objective evidence of in patients with frequent ventricular ectopy without clinically apparent heart disease (Kennedy et al.), 57
Once-a-day treatment of hypertension with pindolol (Frithz), 413
-daily pindolol in hypertension: an ambulatory assessment (Raftery et al.), 417
Open-heart surgery, role of intravenous verapamil in supraventricular tachyarrhythmias after (Gray et al.), 799
Oral administration, chronic, prolongation of verapamil elimination kinetics during (Schwartz et al.), 198
Ostium, aortic, separate, of the left anterior descending and left circumflex coronary arteries from the left aortic sinus of Valsalva (absent left main coronary artery) (Dicicco et al.), 153 (Brief comm.)
Outflow tract tachycardia, exercise provocable right ventricular (Palileo et al.), 185
Oxygen administration and alteration in arterial P_{CO_2}, effects of on ischemic myocardial changes following experimental coronary artery ligation (Weisse et al.), 968

P

Paced cycle length in normals and patients with sinoatrial dysfunction, sinus node recovery time related to (Reiffel et al.), 746
Pacemaker function, ambulatory electrocardiography in the assessment of (Famularo and Kennedy), 1086
prophylactic, embolization, anticoagulation, and long-term success, consideration of in cardioversion of atrial fibrillation (Mancini and Goldberger), 617
Pacing, single-chamber, advantages of dual chamber sequential pacing management of sinus node dysfunction over (Stone, Bhakta, and Lutgen), 1319
transvenous, permanent, symptomatic venous thrombosis of the upper extremity complicating: reversal with streptokinase infusion (Bradof, Sands, and Lakin), 1112 (Brief comm.)
Papillary muscles, stumps of and the mitral anulus, left ventricular incision midway between during mitral valve excision with or without rupture or aneurysmal formation: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
Paraprosthetic mitral regurgitation, severe hemolytic anemia after attempted repair of (Goldberger, Orth, and Moores), 1381 (Brief comm.)
Parotid tumors, carotid sinus syncope as a manifestation of (Mathew, Huang, and Mathew), 316 (Brief comm.)
P_{CO_2}, arterial, alteration in and oxygen administration, effects of on ischemic myocardial changes following experimental coronary artery ligation (Weisse et al.), 968
Pectoris, angina, and cold air exposure (Hirsh), 173 (Letter)
Reply (Feldman and Conti), 173
Penicillin prophylaxis, bacteremia following dental cleaning in patients with and without (Baltch et al.), 1335
Pentoxifylline efficacy in treatment of intermittent claudication: multicenter controlled double-blind trial with objective assessment of chronic occlusive arterial disease patients (Porter et al.), 66
Pericardiocentesis induced intrapericardial thrombus: detection by two-dimensional echocardiography (Schuster and Nanda), 308 (Brief comm.)
Pericarditis, constrictive, massive organized intrapericardial hematoma mimicking (Dunlap et al.), 1373 (Brief comm.)
Peritoneal dialysis, clearance of quinidine during (Hall et al.), 646 (Brief comm.)
Pharmacokinetic(s) comparison of pindolol with other beta-adrenoceptor-blocking agents (Meier), 364
of pindolol in humans and several animal species (Schwarz), 357
Pharmacologic considerations of calcium-channel blockers (Braunwald), 665
differences of beta-adrenoceptor blocking agents: are they relevant? (Clark), 334
Pharmacology, clinical, of adrenergic-adrenoreceptor-blocking drugs (Louis et al.), 407
Phase analysis of radionuclide ventriculograms for the detection of coronary artery disease (Ratib et al.), 1
Phenytoin therapy for ventricular arrhythmias occurring late after surgery for congenital heart disease (Kavey, Blackman, and Sondheimer), 794
Phosphate shift, serum, in acute myocardial infarction (Yaroslavsky et al.), 884 (Letter)
Pindolol, a new beta blocker, effect of on asthma (Cannon, Slavin, and Gonasun), 438
administration, adverse reactions to (Gonasun and Langrall), 482
and methyldopa on blood pressure and plasma norepinephrine, effects of (Winer, Carter, and Eddy), 425
and systolic time intervals in patients with hypertension (Hammond et al.), 456
antihypertensive effects of (Gonasun), 374
clinical pharmacology of (Aellig), 346
:effects on blood pressure and plasma renin activity (Glassock et al.), 421
:experience gained in 10 years of safety monitoring (Krupp and Fanchamps), 486
for ventricular arrhythmia (Podrid and Lown), 491
in hypertension, once-daily: an ambulatory assessment (Raftery et al.), 417
therapeutic trials of: comparison and combination with other drugs (Fanchamps), 388
observations on three dosage forms of (Carruthers), 451
once-a-day treatment of hypertension with (Frithz), 413
pharmacokinetic(s) comparison of with other beta-adrenoceptor-blocking agents (Meier), 364
of in humans and several animal species (Schwarz), 357

Pindolol—*continued*
postmyocardial infarction study: evaluation of the drug's antiarrhythmic and antianginal activity (Welzel et al.), 512
propranolol, and metoprolol, comparison of withdrawal phenomena after (Rangno and Langlois), 473
pulmonary function in hypertensive patients treated with: a report of two studies (Hamilton, Hamilton, and Kirkendall), 432
randomized double-blind study of in patients with stable angina pectoris (Harston and Friesinger), 504
treatment of angina pectoris with: the significance of intrinsic sympathomimetic activity of beta blockers (Kostis et al.), 496
versus methyldopa for hypertension: comparison of adverse reactions (Carr, Mulligan, and Sherrill), 479
Placebo-controlled study in patients with coronary artery spasm: randomized withdrawal from nifedipine (Schick et al.), 690
lack of difference from hydralazine in the long-term treatment of chronic heart failure (Franciosa et al.), 587
Plasma catecholamines and platelet aggregation, in vivo, in acute myocardial infarction (Sorkin et al.), 1255
elevated, without alteration in cardiovascular responsiveness in young men with borderline hypertension (Kawano et al.), 1351
norepinephrine and blood pressure, effects of pindolol and methyldopa on (Winer, Carter, and Eddy), 425
levels, basal, and resting left ventricular performance in patients with congestive heart failure, relationship of exercise capacity to (Francis, Goldsmith, and Cohn), 725
renin activity and blood pressure: pindolol effects on (Glassock et al.), 421
Platelet aggregation and plasma catecholamines, in vivo, in acute myocardial infarction (Sorkin et al.), 1255
Pneumopericardial tamponade postcoronary artery bypass sternal dehiscence (Shawl and Chun), 160 (Brief comm.)
Post-myocardial infarction patients, secondary prevention with beta-adrenoceptor blockers in (Staessen et al.), 1395
study, pindolol: evaluation of the drug's antiarrhythmic and antianginal activity (Welzel et al.), 512
Postoperative abnormalities of interventricular septal motion: two-dimensional and M-mode echocardiographic correlations (Kerber and Litchfield), 263
evaluation of the Fontan procedure by radionuclide angiography (Janos et al.), 785
supraventricular arrhythmias and the role of prophylactic digoxin in cardiac surgery (Chee et al.), 974
Postural hemodynamics and autonomic responses in chronic heart failure, effect of captopril on (Cody), 1190
Potentials, late ventricular, recorded by precordial signal averaging in patients with ventricular tachycardia, effects of antiarrhythmic surgery on (Breithardt et al.), 996
Prazosin, efficacy of low-dose treatment (captopril) and comparison with prazosin: hemodynamic effects of captopril in chronic heart failure (Sharpe and Coxon), 1164
therapy in congestive heart failure (Gould et al.), 1393 (Letter)
Precordial signal averaging in patients with ventricular tachycardia, effects of antiarrhythmic surgery on late ventricular potentials recorded by (Breithardt et al.), 996
Pregnancy, medical treatment of cardiovascular disorders during (Tamari et al.), 1357
Prenatal ultrasound diagnosis of hypoplastic left heart syndrome in utero associated with hydrops fetalis (Sahn et al.), 1368
Pressure, blood, and heart rate response to treadmill exercise, effect of quinidine on (Fenster et al.), 1244
left ventricular end-diastolic, evaluation of changes in by left atrial two-dimensional echocardiography (Haendchen et al.), 740
-volume relations, ventricular diastolic, after the intra-atrial baffle operation for transposition of the great arteries (Sideris et al.), 1045
Primary pulmonary hypertension registry, new (Blaser), 1114 (Letter)
Prinzmetal's angina, ischemic and postischemic ventricular wall motion abnormalities in provoked by hyperventilation (Egeblad, Vilhelmsen, and Mortensen), 1105 (Brief comm.)
Progression, contrasting, of mitral stenosis in Malayans versus American-born Caucasians (Joswig et al.), 1400
Prolapse, mitral valve, left ventricular relaxation, and intracardiac conduction in myotonia atrophica: assessment by digitized echocardiography and noninvasive His bundle recording (Gottdiener et al.), 77
tricuspid valve, detection of (Sasse), 651 (Letter)
Reply (Spodick), 652
Prophylaxis, penicillin, bacteremia following dental cleaning in patients with and without (Baltch et al.), 1335
Propranolol, metoprolol, and pindolol, comparison of withdrawal phenomena after (Rangno and Langlois), 473
nadolol dosage after (Tremblay et al.), 1394 (Letter)
on sinus node function in children, electrophysiologic effects of (Yabek, Berman, and Dillon), 612
Prosthetic heart valve dysfunction, diagnostic value of cinefluoroscopy in the evaluation of (Sands et al.), 622
valve, mitral Björk-Shiley, malfunction of due to septal interference (Jackson, Wolf, and Bloor), 158 (Brief comm.)
Pseudoaneurysm, left ventricular, radionuclide diagnosis of (Gould, Yang, and Martinucci), 1377 (Brief comm.)
Pulmonary artery thrombosis, proximal, noninvasive detection of by two-dimensional echocardiography and computerized tomography (DiCarlo et al.), 879 (Brief comm.)
circulation, effects of acute incremental volume overload on cardiac chamber size, function, and the: analysis by digital intravenous angiography (Slutsky, Carey, and Higgins), 254
disease, chronic obstructive, improvement in cardiac performance by oral long-acting theophylline in (Matthay et al.), 1022
right and left ventricular response to subcutaneous terbutaline in patients with: radionuclide angiographic assessment of cardiac size and function (Hooper et al.), 1027
embolism, time trends in hospital mortality and diagnosis of (Goldhaber and Hennekens), 305 (Brief comm.)
function in hypertensive patients treated with pindolol: a report of two studies (Hamilton, Hamilton, and Kirkendall), 432
hypertension of collagen vascular disease, hemodynamic effects of captopril in (Niarchos et al.), 834
registry, new primary (Blaser), 1114 (Letter)
venous connection, anomalous, total, infradiaphragmatic, in siblings (Baron et al.), 1107 (Brief comm.)
Pulse, finger volume, diminished, in borderline hypertension: evidence for early structural vascular abnormality (Zweifler and Nicholls), 812
rate, blood pressure, and the renin-angiotensin-aldosterone system in patients with essential hypertension, acute effects of the calcium antagonist, nifedipine, on (Hiramatsu et al.), 1346
Purkinje fibers, effects of epinephrine on the electrophysiologic properties of, surviving myocardial infarction (Cameron, Dersham, and Han), 551

Q

Q waves, transient appearance of in coronary disease during exercise electrocardiography: consideration of mechanisms and clinical importance (Bateman et al.), 182
QRS axis, frontal plane, significance of: M-mode echocardiography in left bundle branch block (Strasberg et al.), 775
QT syndrome, the long: effects of drugs and left stellate ganglion block (Milne et al.), 194
Quinidine administration, use of control steady-state serum digoxin levels for predicting serum digoxin concentration after (Friedman and Chen), 72
and other agents on digitalis glycosides, the influence of (Bussey), 289
during peritoneal dialysis, clearance of (Hall et al.), 646 (Brief comm.)
effect of on heart rate and blood pressure response to treadmill exercise (Fenster et al.), 1244

R

Radionuclide angiocardiography, resting and exercise, assessment of left ventricular function by following acute myocardial infarction (Upton et al.), 1232

Radionuclide—*continued*
angiographic assessment of cardiac size and function: right and left ventricular response to subcutaneous terbutaline in patients with chronic obstructive pulmonary disease (Hooper et al.), 1027
evaluation of left ventricular function by resting ejection rate during the first third of systole in patients with chronic aortic regurgitation (Johnson et al.), 92
angiography, diagnosis of tricuspid regurgitation by (Gould, Yang, and Gopalaswamy), 1104 (Brief comm.)
equilibrium, and two-dimensional echocardiography, noninvasive detection of ventricular aneurysm by combined (Sorensen et al.), 145
postoperative evaluation of the Fontan procedure by (Janos et al.), 785
assessment of sequential changes in left and right ventricular function following first acute transmural myocardial infarction (Nemerovski et al.), 709
diagnosis of left ventricular pseudoaneurysm (Gould, Yang, and Martinucci), 1377 (Brief comm.)
ejection fraction and hemodynamic cardiac performance evaluations: patterns of exercise response in patients with severe left ventricular dysfunction (Hecht et al.), 718
evaluation, exercise, combined with hemodynamic evaluation, of left ventricular aneurysmectomy (Dymond et al.), 977
ventriculograms, phase analysis of for the detection of coronary artery disease (Ratib et al.), 1
ventriculography, equilibrium, reproducibility of in patients with coronary artery disease: response of left ventricular ejection fraction and regional wall motion to supine bicycle exercise (Hecht et al.), 567
for assessment of changes in myocardial performance induced by disopyramide phosphate, use of (Kowey et al.), 769
Randomized double-blind study of pindolol in patients with stable angina pectoris (Harston and Friesinger), 504
Recovery time, sinus node, related to paced cycle length in normals and patients with sinoatrial dysfunction (Reiffel et al.), 746
Rectal adenocarcinoma, cardiac metastasis of diagnosed by two-dimensional echocardiography (Henuzet et al.), 637 (Brief comm.)
Reentry, bundle branch, sustained ventricular tachycardia due to (Lloyd et al.), 1095 (Brief comm.)
Refractory angina pectoris, efficacy of nifedipine therapy for (Stone, Turi, and Muller), 672
Regional wall motion and left ventricular ejection fraction, correlation of clinical assessment of: computer-enhanced digital angiography (Vas et al.), 732
response of to supine bicycle exercise: reproducibility of equilibrium radionuclide ventriculography in patients with coronary artery disease (Hecht et al.), 567
Registry, new primary pulmonary hypertension (Blaser), 1114 (Letter)
Regurgitant fraction and right and left ventricular function at rest and during exercise, assessment of: a new technique for determination of right ventricular stroke counts from gated equilibrium blood pool studies (Henze et al.), 953
Regurgitation, aortic, chronic, cardiac response to exercise in patients with (Peter and Jones), 85
mitral, clinically isolated, severe, chronic, pure, etiology of: analysis of 97 patients over 30 years of age having mitral valve replacement (Waller et al.), 276
paraprosthetic, severe hemolytic anemia after attempted repair of (Goldberger, Orth, and Moores), 1381 (Brief comm.)
tricuspid, diagnosis of by radionuclide angiography (Gould, Yang, and Gopalaswamy), 1104 (Brief comm.)
Reinfarction, reocclusion, and recurrent angina following intracoronary thrombolysis with streptokinase in acute myocardial infarction, importance of follow-up medical and surgical approaches to prevent (Lee et al.), 921
Renal, coronary, and systemic hemodynamics in patients with severe congestive heart failure, effect of captopril on (Powers et al.), 1203
Renal—*continued*
dysfunction with sulphinpyrazone therapy (Boelaert et al.), 174 (Letter)
Reply (Warren et al.), 175
Renin activity, plasma, and blood pressure: pindolol effects on (Glassock et al.), 421
-angiotensin-aldosterone activity in the pathophysiology of chronic heart failure, use of captopril to estimate (Cody and Laragh), 1184
system, pulse rate, and blood pressure in patients with essential hypertension, acute effects of the calcium antagonist, nifedipine, on (Hiramatsu et al.), 1346
Reocclusion, reinfarction, and recurrent angina following intracoronary thrombolysis with streptokinase in acute myocardial infarction, importance of follow-up medical and surgical approaches to prevent (Lee et al.), 921
Reperfusion and acute global ischemia, myocardial responses to (Taichman et al.), 1247
Reversibility of bronchospasm in airway obstruction (von Wichert), 446
Right and left ventricular function following first acute transmural myocardial infarction, radionuclide assessment of sequential changes in (Nemerovski et al.), 709
response to subcutaneous terbutaline in patients with chronic obstructive pulmonary disease: radionuclide angiographic assessment of cardiac size and function (Hooper et al.), 1027
aortic sinus, origin of the left main from the right coronary artery or from the with intramyocardial tunneling to the left side of the heart via the ventricular septum: the case against clinical significance of myocardial bridge or coronary tunnel (Roberts et al.), 303 (Brief comm.)
coronary artery, origin of the left main from the or from the right aortic sinus with intramyocardial tunneling to the left side of the heart via the ventricular septum: the case against clinical significance of myocardial bridge or coronary tunnel (Roberts et al.), 303 (Brief comm.)
main coronary artery, fatal atherosclerotic narrowing of the: origin of the left anterior descending or left circumflex coronary artery from the right (the true "left-main equivalent") (Roberts, Waller, and Roberts), 638 (Brief comm.)
ventricular function and size, comparison of single-plane and biplane contrast analyses of (Slutsky et al.), 100
outflow tract tachycardia, exercise provocable (Palileo et al.), 185
Risk(s) and benefits of amiodarone, in perspective (Rotmensch, Belhassen, and Ferguson), 1117
factors of coronary heart disease in black populations. Part II of study (Gillum and Grant), 852
Runners, apparently normal, well-trained, prevalence and nature of cardiac arrhythmias in (Pantano and Oriel), 762

S

Safety monitoring, experience gained in 10 years of—pindolol (Krupp and Fanchamps), 486
Salbutamol, hemodynamic effects of, an oral long-acting beta-stimulant, in patients with congestive heart failure (Mifune et al.), 1011
Saphenous vein associated with early delivery of blood cardioplegia in coronary artery bypass graft surgery, harvesting and protection of the (Feikes and Roberts), 329
Scintigraphic and angiographic techniques, complementary roles of in assessment of the extent of coronary artery disease (Gutman et al.), 653
Second and first heart sounds, frequency analysis approach to the origin of the (Heckman et al.), 1309
Secondary prevention with beta-adrenoceptor blockers in post-myocardial infarction patients (Staessen et al), 1395
Septal aneurysm, ventricular, paradoxically bulging into the left ventricular outflow tract, two-dimensional echocardiographic features of (Barron et al.), 156 (Brief comm.)

Septal—*continued*
hypertrophy, asymmetric, on echocardiographic study, left ventricular thrombus demonstrating canalization and mimicking (Pollick et al.), 641 (Brief comm.)
interference, malfunction of mitral Björk-Shiley prosthetic valve due to (Jackson, Wolf, and Bloor), 158 (Brief comm.)
motion, interventricular, postoperative abnormalities of: two-dimensional and M-mode echocardiographic correlations (Kerber and Litchfield), 263
ventricular, effect of cardiac surgery on: assessment by intraoperative echocardiography and cross-sectional two-dimensional echocardiography (Waggoner et al.), 1271
Sequential coronary angiography, reliability of assessing change with (Detre et al), 816
pacing management, dual chamber, of sinus node dysfunction: advantages over single-chamber pacing (Stone, Bhakta, and Lutgen), 1319
Serum digoxin concentration after quinidine administration, use of control steady-state serum digoxin levels for predicting (Friedman and Chen), 72
levels, control steady-state, use of for predicting serum digoxin concentration after quinidine administration (Friedman and Chen), 72
lidocaine levels, excessive, during maintenance infusions: mechanisms and prevention (Davison, Parker, and Atkinson), 203
phosphate shift in acute myocardial infarction (Yaroslavsky et al.), 884 (Letter)
Siblings, infradiaphragmatic total anomalous pulmonary venous connection in (Baron et al.), 1107 (Brief comm.)
Sick sinus syndrome, sinoatrial arrest due to lidocaine injection in during amiodarone administration (Keidar, Grenadier, and Palant), 1384 (Brief comm.)
Side effects and safety, evaluation of in nifedipine therapy in angina pectoris (Terry), 681
Single-chamber pacing, advantages of dual chamber sequential pacing management of sinus node dysfunction over (Stone, Bhakta, and Lutgen), 1319
-plane and biplane contrast analyses, of right ventricular function and size, comparison of (Slutsky et al.), 100
Sinoatrial arrest due to lidocaine injection in sick sinus syndrome during amiodarone administration (Keidar, Grenadier, and Palant), 1384
dysfunction, sinus node recovery time related to paced cycle length in normals and patients with (Reiffel et al.), 746
Sinus node dysfunction, dual chamber sequential pacing management of: advantages over single-chamber pacing (Stone, Bhakta, and Lutgen), 1319
function in children, electrophysiologic effects of propranolol on (Yabek, Berman, and Dillon), 612
recovery time related to paced cycle length in normals and patients with sinoatrial dysfunction (Reiffel et al.), 746
Sleep apnea, influence of on nocturnal arrhythmias in IHD (Otsuka et al.), 320 (Letter)
Sodium nitroprusside and captopril, comparative analysis of the hemodynamic actions of in severe chronic congestive heart failure (Hermanovich et al.), 1211
Space-occupying lesions of the myocardium: role of two-dimensional echocardiography in detection of cardiac tumors in children (Duncan et al.), 780
Spasm, coronary artery, placebo-controlled study in patients with: randomized withdrawal from nifedipine (Schick et al.), 690
coronary thrombosis, and myocardial infarction: a hypothesis concerning the pathophysiology of acute myocardial infarction (Dalen, Ockene, and Alpert), 1119
ST segment performance of ambulatory ECG monitoring systems (Gallant), 322 (Letter)
Reply (Bragg-Remschel, Anderson, and Winkle), 323
Stable angina pectoris, chronic advantages and disadvantages of combination therapy in: calcium-channel blockers and beta blockers (Krikler, Harris, and Rowland), 702
Steal phenomenon, coronary, demonstration of: left coronary artery-to-left ventricular fistula (Cheng), 870 (Brief comm.)
Stenosis(ses), aortic, sudden death in monitored by ear densitographic pulse and ECG (Nikolic et al.), 311 (Brief comm.)
mitral, effects of nitroglycerin on supine and upright exercise in (Bornheimer et al.), 1288
in Malayans versus American-born Caucasians, contrasting progression of (Joswig et al.), 1400
partial, influence of the autonomic nervous system on coronary blood flow during (Raeder, Verrier, and Lown), 249
severe, of previously confirmed normally functioning aortic valves, documented development of (Roberts et al.), 306 (Brief comm.)
Sternal dehiscence, pneumopericardial tamponade postcoronary artery bypass (Shawl and Chun), 160 (Brief comm.)
Stethoscope, acoustical performance of the: a comparative analysis (Kindig et al.), 269
Stimulation, cardiac, internal, sparkling joules of: cardioversion, defibrillation, and ablation (Fisher, Matos, and Kim), 177
Streptokinase in acute myocardial infarction, intracoronary thrombolysis with, importance of follow-up medical and surgical approaches to prevent reinfarction, reocclusion, and recurrent angina following (Lee et al.), 921
: Western Washington randomized trial-protocol and progress report (Kennedy, Fritz, Ritchie, et al.), 899
infusion, reversal with of symptomatic venous thrombosis of the upper extremity complicating permanent transvenous pacing (Bradof, Sands, and Lakin), 1112 (Brief comm.)
intracoronary, in clinical practice (Timmis et al.), 925
sequential, and transluminal angioplasty in unstable angina with evolving myocardial infarction (Meltzer et al.), 1109 (Brief comm.)
treatment of myocardial infarction with: efficacy and safety data from 209 United States cases in the Hoechst-Roussel registry (Weinstein), 894
intravenous, high-dose, brief, early in acute myocardial infarction (Spann et al.), 939
Stroke counts, right ventricular, a new technique for determination of from gated equilibrium blood pool studies: assessment of regurgitant fraction and right and left ventricular function at rest and during exercise (Henze et al.), 953
Sudden coronary death, definition of (Kayser), 649 (Letter)
Reply (Goldstein), 649
death in aortic stenosis monitored by ear densitographic pulse and ECG (Nikolic et al.), 311 (Brief comm.)
Sulphinpyrazone therapy with renal dysfunction (Boelaert et al.), 174 (Letter)
Reply (Warren et al.), 175
Supine and upright exercise in mitral stenosis, effects of nitroglycerin on (Bornheimer et al.), 1288
Supraventricular arrhythmias, postoperative, and the role of prophylactic digoxin in cardiac surgery (Chee et al.), 974
tachyarrhythmias after open-heart surgery, role of intravenous verapamil in (Gray et al.), 799
tachycardia associated with nodoventricular and concealed atrioventricular bypass tracts (Lerman et al.), 1097 (Brief comm.)
Surgery, cardiac, effect of on ventricular septal motion: assessment by intraoperative echocardiography and cross-sectional two-dimensional echocardiography (Waggoner et al.), 1271
Surgical excision of automatic atrial tachycardia: anatomic and electrophysiologic correlates (Josephson et al.), 1076
Survival and clinical response in patients with congestive heart failure treated with captopril, determinants of (Creager et al.), 1147
Symmetric and asymmetric left ventricular hypertrophy, coexistence of in a family with hypertrophic cardiomyopathy (Ciro, Maron, and Roberts), 643 (Brief comm.)
Sympathetic activity, antihypertensive agents which inhibit: potentially adverse effects of combination treatment (Drayer and Weber), 660
Sympathomimetic activity, intrinsic, is superior to beta-1 selectivity: cardiac safety of acute beta blockade (Heikkilä and Nieminen), 464

Sympathomimetic activity—*continued*
of beta blockers, the significance of intrinsic: treatment of angina pectoris with pindolol (Kostis et al.), 496
Symposium perspective (Mason), 891
Syncope, carotid sinus, as a manifestation of parotid tumors (Mathew, Huang, and Mathew), 316 (Brief comm.)
vasodepressor, circulatory control mechanisms in (Goldstein et al.), 1071
Systemic, renal, and coronary hemodynamics in patients with severe congestive heart failure, effect of captopril on (Powers et al.), 1203
Systolic and diastolic function, left ventricular, in hyperthyroidism (Friedman et al.), 1303
time intervals and echocardiography, relative sensitivity of for assessing acute positive inotropic interventions in normal human subjects (Walsh, Crawford, and O'Rourke), 1061
and pindolol in patients with hypertension (Hammond et al.), 456

T

T waves, hyperacute, revisited (Goldberger), 888
Tachyarrhythmias, supraventricular, after open-heart surgery, role of intravenous verapamil in (Gray et al.), 799
Tachycardia, automatic atrial, surgical excision of: anatomic and electrophysiologic correlates (Josephson et al.), 1076
exercise provocable right ventricular outflow tract (Palileo et al.), 185
recurrent ventricular, additional observations on transvenous cardioversion of (Zipes et al.), 163 (Brief comm.)
supraventricular, associated with nodoventricular and concealed atrioventricular bypass tracts (Lerman et al.), 1097 (Brief comm.)
ventricular, effects of antiarrhythmic surgery on late ventricular potentials recorded by precordial signal averaging in patients with (Breithardt et al.), 996
electrocardiographic changes following endocardial resection for (Kienzle et al.), 753
macroreentrant, sustained (Welch et al.), 166 (Brief comm.)
prediction of response to class I antiarrhythmic drugs during electrophysiologic study of (Swiryn et al.), 43
sustained, due to bundle branch reentry (Lloyd et al.), 1095 (Brief comm.)
long-term results of endocardial resection for in coronary disease patients (Josephson, Harken, and Horowitz), 51
Takayasu's aortitis, etiologic aspects of coagulopathy in (Kanaide, Takeshita, and Nakamura), 1039
Tamponade, pneumopericardial, postcoronary artery bypass sternal dehiscence (Shawl and Chun), 160 (Brief comm.)
Terbutaline and captopril therapy, complementary combined, in severe chronic congestive heart failure (Awan et al.), 1224
intravenous, beneficial hemodynamic effects of in patients with severe heart failure (Wang et al.), 1016
subcutaneous, right and left ventricular response to in patients with chronic obstructive pulmonary disease:radionuclide angiographic assessment of cardiac size and function (Hooper et al.), 1027
Theophylline, oral long-acting, improvement in cardiac performance by in chronic obstructive pulmonary disease (Matthay et al.), 1022
Therapeutic trials of pindolol in hypertension: comparison and combination with other drugs (Fanchamps), 388
Therapy of severe chronic congestive heart failure (Awan and Massie), 1125
Thoracic impedance cardiography (Krohn), 1392 (Letter)
(Spodick), 1392
Reply (Sheps), 1392
Three dosage forms of pindolol, observations on (Carruthers), 451
Thrombolysis, intracoronary, in syndromes of unstable ischemia: angiographic and clinical results (Vetrovec et al.), 946
up to 18 hours after onset of pain in evolving myocardial infarction (Smalling et al.), 912
with streptokinase in acute myocardial infarction, importance of follow-up medical and surgical approaches to prevent reinfarction, reocclusion, and recurrent angina following (Lee et al.), 921
Thrombosis, coronary, coronary spasm, and myocardial infarction: a hypothesis concerning the pathophysiology of acute myocardial infarction (Dalen, Ockene, and Alpert), 1119
proximal pulmonary artery, noninvasive detection of by two-dimensional echocardiography and computerized tomography (DiCarlo et al.), 879 (Brief comm.)
venous, symptomatic, of the upper extremity complicating permanent transvenous pacing: reversal with streptokinase infusion (Bradof, Sands, and Lakin), 1112 (Brief comm.)
Thromboxane concentrations, influence of blood sampling site and technique on in patients with ischemic heart disease (Hirsh et al.), 234
Thrombus, intrapericardial, pericardiocentesis induced: detection by two-dimensional echocardiography (Schuster and Nanda), 308 (Brief comm.)
left ventricular, demonstrating canalization and mimicking asymmetric septal hypertrophy on echocardiographic study (Pollick et al.), 641 (Brief comm.)
Time intervals, systolic, and pindolol in patients with hypertension (Hammond et al.), 456
trends in hospital mortality and diagnosis of pulmonary embolism (Goldhaber and Hennekens), 305 (Brief comm.)
Tomography, computerized, and two-dimensional echocardiography, noninvasive detection of proximal pulmonary artery thrombosis by (DiCarlo et al.) 879 (Brief comm.)
Toxicity, digitalis, transient atrial dissociation due to (Condorelli, Dato, and Sindoni), 1102 (Brief comm.)
Transient atrial dissociation due to digitalis toxicity (Condorelli, Dato, and Sindoni), 1102 (Brief comm.)
Transmural myocardial infarction, comparison of clinical signs and hemodynamic state in the early hours of (Shell et al.), 521
first acute, radionuclide assessment of sequential changes in left and right ventricular function following (Nemerovski et al.), 709
infarct size, relation of healed to length of survival after acute myocardial infarction, age at death, and amount and extent of coronary arterial narrowing by atherosclerotic plaques: analysis of 70 necropsy patients (Cabin and Roberts), 216
Transposition of the great arteries, corrected, two-dimensional echocardiographic identification of high anatomically inserted venous atrioventricular valve in the diagnosis of (Motro et al.), 1376 (Brief comm.)
ventricular diastolic pressure-volume relations after the intra-atrial baffle operation for (Sideris et al.), 1045
Transvenous cardioversion of recurrent ventricular tachycardia, additional observations on (Zipes et al.), 163 (Brief comm.)
pacing, permanent, symptomatic venous thrombosis of the upper extremity complicating: reversal with streptokinase infusion (Bradof, Sands, and Lakin), 1112 (Brief comm.)
Treadmill exercise, effect of quinidine on the heart rate and blood pressure response to (Fenster et al.), 1244
protocols predischarge, selected, and six weeks postinfarction, superiority of for detecting ischemic abnormalities (Starling, Crawford, and O'Rourke), 1054
test performance, exercise, patterns of in patients with left main coronary artery disease: detection dependent on left coronary dominance or coexistent dominant right coronary disease (Stone, LaFollette, and Cohn), 13
Tricuspid regurgitation, diagnosis of by radionuclide angiography (Gould, Yang, and Gopalaswamy), 1104 (Brief comm.)
valve prolapse, detection of (Sasse), 651 (Letter)
Reply (Spodick), 652
Tumors, cardiac, in children, role of two-dimensional echocardiography in detection of: space-occupying lesions of the myocardium (Duncan et al.), 780
parotid, carotid sinus syncope as a manifestation of (Mathew, Huang, and Mathew), 316 (Brief comm.)
Tunneling, intramyocardial, origin of the left main from the right coronary artery or from the right aortic sinus with to the left side of the heart via the ventricular septum: the case against clinical significance of myocardial bridge or coronary tunnel (Roberts et al.), 303 (Brief comm.)

Two-dimensional and M-mode echocardiographic correlations: postoperative abnormalities of interventricular septal motion (Kerber and Litchfield), 263
features of left ventricular myxoma (Mazer and Harrigan), 875 (Brief comm.)
echocardiographic features of coronary arteriovenous fistula (Rodgers et al.), 872 (Brief comm.)
of ventricular septal aneurysm paradoxically bulging into the left ventricular outflow tract (Barron et al.), 156 (Brief comm.)
identification of high anatomically inserted venous atrioventricular valve in the diagnosis of corrected transposition of the great arteries (Motro et al.), 1376 (Brief comm.)
measurement of left ventricular ejection fraction: prospective analysis of what constitutes an adequate determination (Stamm et al.), 136
echocardiography and computerized tomography, noninvasive detection of proximal pulmonary artery thrombosis by (DiCarlo et al.), 879 (Brief comm.)
and equilibrium radionuclide angiography, noninvasive detection of ventricular aneurysm by combined (Sorensen et al.), 145
cardiac metastasis of rectal adenocarcinoma diagnosed by (Henuzet et al.), 637 (Brief comm.)
cross-sectional, and intraoperative echocardiography, assessment of effect of cardiac surgery on ventricular septal motion by (Waggoner et al.), 1271
detection of atrioventricular disc valve malfunction by (Mehlman and Talano), 1378 (Brief comm.)
early diagnosis of nontransmural myocardial infarction by (Loh et al.), 963
in detection of cardiac tumors in children, role of: space-occupying lesions of the myocardium (Duncan et al.), 780
left atrial, evaluation of changes in left ventricular end-diastolic pressure by (Haendchen et al.), 740
real-time, determination of left ventricular ejection fraction by visual estimation during (Rich et al.), 603

U

Ultrasound diagnosis, prenatal, of hypoplastic left heart syndrome in utero associated with hydrops fetalis (Sahn et al.), 1368
Unstable angina, inpatient treatment of: clinical perspective and sequential management (Helfant), 697
ischemia, intracoronary thrombolysis in syndromes of: angiographic and clinical results (Vetrovec et al.), 946
Upright and supine exercise in mitral stenosis, effects of nitroglycerin on (Bornheimer et al.), 1288
Urinary catecholamines, cardiac rhythm, and arterial pressure in borderline and established essential hypertension, diurnal variations of (Messerli et al.), 109

V

Valsalva, left aortic sinus of, separate aortic ostium of the left anterior descending and left circumflex coronary arteries from the (absent left main coronary artery) (Dicicco et al.), 153 (Brief comm.)
Valve(s), aortic, normally functioning, previously confirmed, documented development of severe stenoses of (Roberts et al.), 306 (Brief comm.)
atrioventricular, high anatomically inserted venous, two-dimensional echocardiographic identification of in the diagnosis of corrected transposition of the great arteries (Motro et al.), 1376 (Brief comm.)
disease, mitral, correction of without valve replacement (Bonchek), 865
dysfunction, heart, prosthetic, diagnostic value of cinefluoroscopy in the evaluation of (Sands et al.), 622
excision, mitral, with or without rupture or aneurysmal formation, left ventricular incision midway between the mitral anulus and stumps of the papillary muscles during: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
malfunction, disc, atrioventricular, detection of by two-dimensional echocardiography (Mehlman and Talano), 1378 (Brief comm.)
Valve—*continued*
prolapse, tricuspid, detection of (Sasse), 651 (Letter) Reply (Spodick), 652
prosthetic, mitral Björk-Shiley, malfunction of due to septal interference (Jackson, Wolf, and Bloor), 158 (Brief comm.)
replacement, complication of active infective endocarditis of the aortic valve with ring abscess treated by: acquired cor triatriatum (left ventricular false aneurysm) (McManus et al.), 312 (Brief comm.)
correction of mitral valve disease without (Bonchek), 865
mitral, analysis of 97 patients over 30 years of age having: etiology of clinically isolated, severe, chronic, pure mitral regurgitation (Waller et al.), 276
Variant angina, regional coronary artery dilation response in (Hill et al.), 226
Vascular abnormality, early structural, evidence for: diminished finger volume pulse in borderline hypertension (Zweifler and Nicholls), 812
disease, collagen, hemodynamic effects of captopril in pulmonary hypertension of (Niarchos et al.), 834
Vasodepressor syncope, circulatory control mechanisms in (Goldstein, et al.), 1071
Vasodilator drugs, other, and captopril, comparison of: contrasting hemodynamic responses in severe heart failure (Packer, Medina, and Yushak), 1215
therapy of severe congestive heart failure: the special importance of angiotensin-converting enzyme inhibition with captopril (Awan and Mason), 1127
Venous atrioventricular valve, high anatomically inserted, two-dimensional echocardiographic identification of in the diagnosis of corrected transposition of the great arteries (Motro et al.), 1376 (Brief comm.)
connection, pulmonary, infradiaphragmatic total anomalous, in siblings (Baron et al.), 1107 (Brief comm.)
thrombosis, symptomatic, of the upper extremity complicating permanent transvenous pacing: reversal with streptokinase infusion (Bradof, Sands, and Lakin), 1112 (Brief comm.)
Ventricular aneurysm, noninvasive detection of by combined two-dimensional echocardiography and equilibrium radionuclide angiography (Sorensen et al.), 145
arrhythmias and regional myocardial blood flow following one-stage and two-stage coronary artery occlusion in anesthetized dogs (Kabell et al.), 537
comparison of two methods of analyzing frequency of (Shapiro et al.), 874 (Brief comm.)
electrophysiologic effects of chronic amiodarone therapy in patients with (Finerman et al.), 987
occurring late after surgery for congenital heart disease, phenytoin therapy for (Kavey, Blackman, and Sondheimer), 794
pindolol for (Podrid and Lown), 491
depolarizations, concealed, due to amiodarone, electrophysiologic observations of (Ezri et al.), 169 (Brief comm.)
diastolic pressure-volume relations after the intra-atrial baffle operation for transposition of the great arteries (Sideris et al.), 1045
ectopy, frequent, without clinically apparent heart disease, objective evidence of occult myocardial dysfunction in patients with (Kennedy et al.), 57
fibrillation complicating acute myocardial infarction: reasons against the routine use of lidocaine (Carruth and Silverman), 545
function and left and right ventricular volumes in chronic heart failure, acute and long-term effects of captopril on (Massie, Kramer, and Topic), 1197
left and right, following first acute transmural myocardial infarction, radionuclide assessment of sequential changes in (Nemerovski et al.), 709
right and left, at rest and during exercise, assessment of regurgitant fraction and: a new technique for determination of right ventricular stroke counts from gated equilibrium blood pool studies (Henze et al.), 953
incision, left, midway between the mitral anulus and the stumps of the papillary muscles during mitral valve excision with or without rupture or aneurysmal formation: analysis of 10 necropsy patients (Roberts, Isner, and Virmani), 1278
performance and myocardial perfusion following exercise training, noninvasive assessment of changes in (Tubau et al.), 238

Ventricular—*continued*
septal motion, effect of cardiac surgery on: assessment by intraoperative echocardiography and cross-sectional two-dimensional echocardiography (Waggoner et al.), 1271
septum, origin of the left main from the right coronary artery or from the right aortic sinus with intramyocardial tunneling to the left side of the heart via the: the case against clinical significance of myocardial bridge or coronary tunnel (Roberts et al.), 303 (Brief comm.)
tachycardia, effects of antiarrhythmic surgery on late ventricular potentials recorded by precordial signal averaging in patients with (Breithardt et al.), 996
electrocardiographic changes following endocardial resection for (Kienzle et al.), 753
macroreentrant, sustained (Welch et al.), 166 (Brief comm.)
prediction of response to class I antiarrhythmic drugs during electrophysiologic study of (Swiryn et al.), 43
recurrent, additional observations on transvenous cardioversion of (Zipes et al.), 163 (Brief comm.)
sustained, due to bundle branch reentry (Lloyd et al.), 1095 (Brief comm.)
long-term results of endocardial resection for in coronary disease patients (Josephson, Harken, and Horowitz), 51
volumes, left and right, and ventricular function in chronic heart failure, acute and long-term effects of captopril on (Massie, Kramer, and Topic), 1197
wall motion abnormalities, ischemic and postischemic, in Prinzmetal's angina provoked by hyperventilation (Egeblad, Vilhelmsen, and Mortensen), 1105 (Brief comm.)
Ventriculograms, left, and body surface isopotential maps, correlation between in patients with old inferoposterior myocardial infarction (Ohta et al.), 1262
radionuclide, phase analysis of for the detection of coronary artery disease (Ratib et al.), 1
Ventriculography, equilibrium radionuclide, reproducibility of in patients with coronary artery disease: response of left ventricular ejection fraction and regional wall motion to supine bicycle exercise (Hecht et al.), 567
Ventriculography—*continued*
radionuclide, for assessment of changes in myocardial performance induced by disopyramide phosphate, use of (Kowey et al.), 769
Verapamil elimination kinetics, prolongation of during chronic oral administration (Schwartz et al.), 198
intravenous, role of in supraventricular tachyarrhythmias after open-heart surgery (Gray et al.), 799
therapy, rapid atrial fibrillation during hypercalcemia despite (Santo, Aderka, and Pinkhas), 320 (Letter)
Viral hemorrhagic fevers, myocardial damage in (Milei and Bolomo), 1385 (Brief comm.)
Volume overload, incremental, acute, effects of on cardiac chamber size, function, and the pulmonary circulation: analysis by digital intravenous angiography (Slutsky, Carey, and Higgins), 254

W

Wall motion abnormalities, ventricular, ischemic and postischemic, in Prinzmetal's angina provoked by hyperventilation (Egeblad, Vilhelmsen, and Mortensen), 1105 (Brief comm.)
Washington, Western, randomized trial-protocol and progress report: streptokinase in acute myocardial infarction (Kennedy, Fritz, Ritchie, et al.), 899
Waves, atrial fibrillation, in the geriatric population, clinicopathologic study on determinants of the amplitude of (Inoue et al.), 1382 (Brief comm.)
T, hyperacute, revisited (Goldberger), 888
Western Washington randomized trial-protocol and progress report: streptokinase in acute myocardial infarction (Kennedy, Fritz, Ritchie, et al.), 899
Wilson's disease, fatal cardiac complications of (Kuan), 314 (Brief comm.)
Withdrawal from nifedipine, randomized: placebo-controlled study in patients with coronary artery spasm (Schick et al.), 690
of beta-blocking drugs (Walden et al.), 515
phenomena after propranolol, metoprolol, and pindolol, comparison of (Rangno and Langlois), 473